

Medicinal plants

Medicinal plants, also called **medicinal herbs**, have been discovered and used in traditional medicine practices since prehistoric times. Plants synthesise hundreds of chemical compounds for functions including defence against insects, fungi, diseases, and herbivorous mammals. Numerous phytochemicals with potential or established biological activity have been identified. However, since a single plant contains widely diverse phytochemicals, the effects of using a whole plant as medicine are uncertain. Further, the phytochemical content and pharmacological actions, if any, of many plants having medicinal potential remain unassessed by rigorous scientific research to define efficacy and safety.^[2]

The earliest historical records of herbs are found from the Sumerian civilisation, where hundreds of medicinal plants including opium are listed on clay tablets. The Ebers Papyrus from ancient Egypt, c. 1550 BC, describes over 850 plant medicines. The Greek physician Dioscorides, who worked in the Roman army, documented over 1000 recipes for medicines using over 600 medicinal plants in *De materia medica*, c. 60 AD; this formed the basis of pharmacopoeias for some 1500 years. Drug research makes use of ethnobotany to search for pharmacologically active substances in nature, and has in this way discovered hundreds of useful compounds. These include the common drugs aspirin, digoxin, quinine, and opium. The compounds found in plants are of many kinds, but most are in four major biochemical classes: alkaloids, glycosides, polyphenols, and terpenes.

Medicinal plants are widely used in non-industrialized societies, mainly because they are readily available and cheaper than modern medicines. The annual global export value of the thousands of types of plants with suspected medicinal properties was estimated to be US\$2.2 billion in 2012.^[3] In 2017, the potential global market for botanical extracts and medicines was estimated at several hundred billion dollars.^[2] In many countries, there is little regulation of traditional medicine, but the World Health Organization coordinates a network to encourage safe and rational usage. Medicinal plants face both general threats, such as climate change and habitat destruction, and the specific threat of over-collection to meet market demand.^[2]



The bark of willow trees contains salicylic acid, the active metabolite of aspirin, and has been used for millennia to relieve pain and reduce fever.^[1]



Medicinal plants

Contents

History

Prehistoric times

Ancient times

Middle Ages
Early Modern
19th and 20th centuries

Context

Phytochemical basis

Alkaloids
Glycosides
Polyphenols
Terpenes

In practice

Cultivation
Preparation
Usage
Effectiveness
Regulation
Drug discovery
Safety
Quality, advertising, and labelling

Threats

See also

Notes

References

Further reading

History

Prehistoric times

Plants, including many now used as culinary herbs and spices, have been used as medicines, not necessarily effectively, from prehistoric times. Spices have been used partly to counter food spoilage bacteria, especially in hot climates,^{[5][6]} and especially in meat dishes which spoil more readily.^[7] Angiosperms (flowering plants) were the original source of most plant medicines.^[8] Human settlements are often surrounded by weeds used as herbal medicines, such as nettle, dandelion and chickweed.^{[9][10]} Humans were not alone in using herbs as medicines: some animals such as non-human primates, monarch butterflies and sheep ingest medicinal plants when they are ill.^[11] Plant samples from prehistoric burial sites are among the lines of evidence that Paleolithic peoples had knowledge of herbal medicine. For instance, a 60 000-year-old Neanderthal burial site, "Shanidar IV", in northern Iraq has yielded large amounts of pollen from eight plant species, seven of which are used now as herbal remedies.^[12] A mushroom was found in the personal effects of Ötzi the Iceman, whose body was frozen in the Ötztal Alps for more than 5,000 years. The mushroom was probably used against whipworm.^[13]



Dioscorides's 1st century *De materia medica*, seen here in a c. 1334 copy in Arabic, describes some 1000 drug recipes based on over 600 plants.^[4]

Ancient times

In ancient Sumeria, hundreds of medicinal plants including myrrh and opium are listed on clay tablets. The ancient Egyptian Ebers Papyrus lists over 800 plant medicines such as aloe, cannabis, castor bean, garlic, juniper, and mandrake.^[14] From ancient times to the present, Ayurvedic medicine as documented in the Atharva Veda, the Rig Veda and the Sushruta Samhita has used hundreds of pharmacologically active herbs and spices such as turmeric, which contains curcumin.^{[15][16]} The Chinese pharmacopoeia, the Shennong Ben Cao Jing records plant medicines such as chaulmoogra for leprosy, ephedra, and hemp.^[17] This was expanded in the Tang Dynasty Yaoxing Lun.^[18] In the fourth century BC, Aristotle's pupil Theophrastus wrote the first systematic botany text, Historia plantarum.^[19] In around 60 AD, the Greek physician Pedanius Dioscorides, working for the Roman army, documented over 1000 recipes for medicines using over 600 medicinal plants in De materia medica. The book remained the authoritative reference on herbalism for over 1500 years, into the seventeenth century.^[4]



The Ebers Papyrus (c. 1550 BC) from Ancient Egypt describes the use of hundreds of plant medicines.^[14]

Middle Ages

In the Early Middle Ages, Benedictine monasteries preserved medical knowledge in Europe, translating and copying classical texts and maintaining herb gardens.^{[20][21]} Hildegard of Bingen wrote *Causae et Curae* ("Causes and Cures") on medicine.^[22] In the Islamic Golden Age, scholars translated many classical Greek texts including Dioscorides into Arabic, adding their own commentaries.^[23] Herbalism flourished in the Islamic world, particularly in Baghdad and in Al-Andalus. Among many works on medicinal plants, Abulcasis (936–1013) of Cordoba wrote *The Book of Simples*, and Ibn al-Baitar (1197–1248) recorded hundreds of medicinal herbs such as Aconitum, nux vomica, and tamarind in his *Corpus of Simples*.^[24] Avicenna included many plants in his 1025 *The Canon of Medicine*.^[25] Abu-Rayhan Biruni,^[26] Ibn Zuhr,^[27] Peter of Spain, and John of St Amand wrote further pharmacopoeias.^[28]



Illustration of a 1632 copy of Avicenna's 1025 *The Canon of Medicine*, showing a physician talking to a female patient in a garden, while servants prepare medicines.

Early Modern

The Early Modern period saw the flourishing of illustrated herbals across Europe, starting with the 1526 *Grete Herball*. John Gerard wrote his famous *The Herball or General History of Plants* in 1597, based on Rembert Dodoens, and Nicholas Culpeper published his *The English Physician Enlarged*.^[29] Many new plant medicines arrived in Europe as products of Early Modern exploration and the resulting Columbian Exchange, in which livestock, crops and technologies were transferred between the Old World and the Americas in the 15th and 16th centuries. Medicinal herbs arriving in the Americas included garlic, ginger, and turmeric; coffee, tobacco and coca travelled in the other direction.^{[30][31]} In Mexico, the sixteenth century *Badianus Manuscript* described medicinal plants available in Central America.^[32]

19th and 20th centuries

The place of plants in medicine was radically altered in the 19th century by the application of chemical analysis. Alkaloids were isolated from a succession of medicinal plants, starting with morphine from the poppy in 1806, and soon followed by ippecacuanha and strychnos in 1817, quinine from the cinchona tree, and then many others. As chemistry progressed, additional classes of pharmacologically active substances were discovered in medicinal plants.^{[33][34]} Commercial extraction of purified alkaloids including morphine from medicinal plants began at Merck in 1826. Synthesis of a substance first discovered in a medicinal plant began with salicylic acid in 1853.^[34] Around the end of the 19th century, the mood of pharmacy turned against medicinal plants, as enzymes often modified the active ingredients when whole plants were dried, and alkaloids and glycosides purified from plant material started to be preferred.^[33] Drug discovery from plants continued to be important through the 20th century and into the 21st, with important anti-cancer drugs from yew and Madagascar periwinkle.^[34]



An early illustrated book of medicinal plants,^[29] The Grete Herball, 1526

Context

Medicinal plants are used with the intention of maintaining health, to be administered for a specific condition, or both, whether in modern medicine or in traditional medicine.^{[2][35]} The Food and Agriculture Organization estimated in 2002 that over 50,000 medicinal plants are used across the world.^[36] The Royal Botanic Gardens, Kew more conservatively estimated in 2016 that 17,810 plant species have a medicinal use, out of some 30,000 plants for which a use of any kind is documented.^[37]

In modern medicine, around a quarter^[a] of the drugs prescribed to patients are derived from medicinal plants, and they are rigorously tested.^{[35][38]} In other systems of medicine, medicinal plants may constitute the majority of what are often informal attempted treatments, not tested scientifically.^[39] The World Health Organization estimates, without reliable data, that some 80 percent of the world's population depends mainly on traditional medicine (including but not limited to plants); perhaps some two billion people are largely reliant on medicinal plants.^{[35][38]} The use of plant-based materials including herbal or natural health products with supposed health benefits, is increasing in developed countries.^[40] This brings attendant risks of toxicity and other effects on human health, despite the safe image of herbal remedies.^[40] Herbal medicines have been in use since long before modern medicine existed; there was and often still is little or no knowledge of the pharmacological basis of their actions, if any, or of their safety. The World Health Organization formulated a policy on traditional medicine in 1991, and since then has published guidelines for them, with a series of monographs on widely used herbal medicines.^{[41][42]}

Medicinal plants may provide three main kinds of benefit: health benefits to the people who consume them as medicines; financial benefits to people who harvest, process, and distribute them for sale; and society-wide benefits, such as job opportunities, taxation income, and a healthier labour force.^[35] However, development of plants or extracts having potential medicinal uses is blunted by weak scientific evidence, poor practices in the process of drug development, and insufficient financing.^[2]

Phytochemical basis


All plants produce chemical compounds which give them an evolutionary advantage, such as defending against herbivores or, in the example of salicylic acid, as a hormone in plant defenses.^{[43][44]} These phytochemicals have potential for use as drugs, and the content and known pharmacological activity of these substances in medicinal plants is the scientific basis for their use in modern medicine, if scientifically confirmed.^[2] For instance, daffodils (Narcissus) contain nine groups of alkaloids including galantamine,

licensed for use against Alzheimer's disease. The alkaloids are bitter-tasting and toxic, and concentrated in the parts of the plant such as the stem most likely to be eaten by herbivores; they may also protect against parasites.^{[45][46][47]}

Modern knowledge of medicinal plants is being systematised in the Medicinal Plant Transcriptomics Database, which by 2011 provided a sequence reference for the transcriptome of some thirty species.^[48] The major classes of pharmacologically active phytochemicals are described below, with examples of medicinal plants that contain them.^{[8][42][49][50][51]}

Alkaloids

Alkaloids are bitter-tasting chemicals, very widespread in nature, and often toxic, found in many medicinal plants.^[52] There are several classes with different modes of action as drugs, both recreational and pharmaceutical. Medicines of different classes include atropine, scopolamine, and hyoscyamine (all from nightshade),^[53] the traditional medicine berberine (from plants such as Berberis and Mahonia),^[b] caffeine (Coffea), cocaine (Coca), ephedrine (Ephedra), morphine (opium poppy), nicotine (tobacco),^[c] reserpine (Rauvolfia serpentina), quinidine and quinine (Cinchona), vincamine (Vinca minor), and vincristine (Catharanthus roseus).^{[51][56]}



The opium poppy Papaver somniferum is the source of the alkaloids morphine and codeine.^[51]

The alkaloid nicotine from tobacco binds directly to the body's Nicotinic acetylcholine receptors, accounting for its pharmacological effects.^[57]

Deadly nightshade, Atropa belladonna, yields tropane alkaloids including atropine, scopolamine and hyoscyamine.^[53]

Glycosides

Anthraquinone glycosides are found in medicinal plants such as rhubarb, cascara, and Alexandrian senna.^{[58][59]} Plant-based laxatives made from such plants include senna,^[60] rhubarb^[61] and Aloe.^[51]


The cardiac glycosides are powerful drugs from medicinal plants including foxglove and lily of the valley. They include digoxin and digitoxin which support the beating of the heart, and act as diuretics.^[43]



Senna alexandrina, containing anthraquinone glycosides, has been used as a laxative for millennia.^[60]



The foxglove, Digitalis purpurea, contains digoxin, a cardiac glycoside. The plant was used on heart conditions long before the glycoside was identified.^{[43][62]}




Digoxin is used to treat atrial fibrillation, atrial flutter and sometimes heart failure.^[43]

Polyphenols

Polyphenols of several classes are widespread in plants, having diverse roles in defenses against plant diseases and predators.^[43] They include hormone-mimicking phytoestrogens and astringent tannins.^{[51][63]} Plants containing phytoestrogens have been administered for centuries for gynecological disorders, such as fertility, menstrual, and menopausal problems.^[64] Among these plants are Pueraria mirifica,^[65] kudzu,^[66] angelica,^[67] fennel, and anise.^[68]


Many polyphenolic extracts, such as from grape seeds, olives or maritime pine bark, are sold as dietary supplements and cosmetics without proof or legal health claims for beneficial health effects.^[69] In Ayurveda, the astringent rind of the pomegranate, containing polyphenols called punicalagins, is used as a medicine.^[70]



Angelica, containing Polyphenols include phytoestrogens (top and middle), mimics of animal gynaecological disorders. estrogen (bottom).^[71]

Terpenes

Terpenes and terpenoids of many kinds are found in a variety of medicinal plants,^[72] and in resinous plants such as the conifers. They are strongly aromatic and serve to repel herbivores. Their scent makes them useful in essential oils, whether for perfumes such as rose and lavender, or for aromatherapy.^{[51][73][74]} Some have medicinal uses: for example, thymol is an antiseptic and was once used as a vermifuge (anti-worm medicine).^[75]



The essential oil of common thyme (*Thymus vulgaris*), contains the monoterpene thymol, an antiseptic and antifungal.^[75]

Thymol is one of many terpenes found in plants.^[75]

In practice

Cultivation

Medicinal plants demand intensive management. Different species each require their own distinct conditions of cultivation. The World Health Organization recommends the use of rotation to minimise problems with pests and plant diseases. Cultivation may be traditional or may make use of conservation agriculture practices to maintain organic matter in the soil and to conserve water, for example with no-till farming systems.^[76] In many medicinal and aromatic plants, plant characteristics vary widely with soil type and cropping strategy, so care is required to obtain satisfactory yields.^[77]



Licensed commercial cultivation of opium poppies, Tasmania, 2010

Preparation

Medicinal plants are often tough and fibrous, requiring some form of preparation to make them convenient to administer. According to the Institute for Traditional Medicine, common methods for the preparation of herbal medicines include decoction, powdering, and extraction with alcohol, in each case yielding a mixture of substances. Decoction involves crushing and then boiling the plant material in water to produce a liquid extract that can be taken orally or applied topically.^[78] Powdering involves drying the plant material and then crushing it to yield a powder that can be compressed into tablets. Alcohol extraction involves soaking the plant material in cold wine or distilled spirit to form a tincture.^[79]



A Medieval physician preparing an extract from a medicinal plant, from an Arabic Dioscorides, 1224

Traditional poultices were made by boiling medicinal plants, wrapping them in a cloth, and applying the resulting parcel externally to the affected part of the body.^[80]

When modern medicine has identified a drug in a medicinal plant, commercial quantities of the drug may either be synthesised or extracted from plant material, yielding a pure chemical.^[34] Extraction can be practical when the compound in question is complex.^[81]

Usage

Plant medicines are in wide use around the world.^[82] In most of the developing world, especially in rural areas, local traditional medicine, including herbalism, is the only source of health care for people, while in the developed world, alternative medicine including use of dietary supplements is marketed aggressively using the claims of traditional medicine. As of 2015, most products made from medicinal plants had not been tested for their safety and efficacy, and products that were marketed in developed economies and provided in the undeveloped world by traditional healers were of uneven quality, sometimes containing dangerous contaminants.^[83] Traditional Chinese medicine makes use of a wide variety of plants, among other materials and techniques.^[84] Researchers from Kew Gardens found 104 species used for diabetes in Central America, of which seven had been identified in at least three separate studies.^{[85][86]} The Yanomami of the Brazilian Amazon, assisted by researchers, have described 101 plant species used for traditional medicines.^{[87][88]}



A herbalist's shop in the souk of Marrakesh, Morocco

Drugs derived from plants including opiates, cocaine and cannabis have both medical and recreational uses. Different countries have at various times made use of illegal drugs, partly on the basis of the risks involved in taking psychoactive drugs.^[89]

Effectiveness

Plant medicines have often not been tested systematically, but have come into use informally over the centuries. By 2007, clinical trials had demonstrated potentially useful activity in nearly 16% of herbal medicines; there was limited in vitro or in vivo evidence for roughly half the medicines; there was only phytochemical evidence for around 20%; 0.5% were allergenic or toxic; and some 12% had basically never been studied scientifically.^[42] Cancer Research UK caution that there is no reliable evidence for the effectiveness of herbal remedies for cancer.^[90]

A 2012 phylogenetic study built a family tree down to genus level using 20,000 species to compare the medicinal plants of three regions, Nepal, New Zealand and the South African Cape. It discovered that the species used traditionally to treat the same types of condition belonged to the same groups of plants in all three regions, giving a "strong phylogenetic signal".^[91] Since many plants that yield pharmaceutical drugs belong to just these groups, and the groups were independently used in three different world regions, the results were taken to mean 1) that these plant groups do have potential for medicinal efficacy, 2) that undefined pharmacological activity is associated with use in traditional medicine, and 3) that the use of a phylogenetic groups for medicines in one region may predict their use in the other regions.^[91]



The bark of the cinchona tree contains the alkaloid quinine, traditionally given for malaria.

Regulation

The World Health Organization (WHO) has been coordinating a network called the International Regulatory Cooperation for Herbal Medicines to try to improve the quality of medical products made from medicinal plants and the claims made for them.^[92] In 2015, only around 20% of countries had well-functioning



The practice of Ayurveda in India, such as the running of this Ayurvedic pharmacy in Rishikesh, is regulated by a government department, AYUSH.

regulatory agencies, while 30% had none, and around half had limited regulatory capacity.^[83] In India, where Ayurveda has been practised for centuries, herbal remedies are the responsibility of a government department, AYUSH, under the Ministry of Health & Family Welfare.^[93]

WHO has set out a strategy for traditional medicines^[94] with four objectives: to integrate them as policy into national healthcare systems; to provide knowledge and guidance on their safety, efficacy, and quality; to increase their availability and affordability; and to promote their rational, therapeutically sound usage.^[94] WHO notes in the strategy that countries are experiencing seven challenges to such implementation, namely in developing and enforcing policy; in integration; in safety and quality, especially in assessment of products and qualification of practitioners; in controlling advertising; in research and development; in education and training; and in the

sharing of information.^[94]

Drug discovery

The pharmaceutical industry has roots in the apothecary shops of Europe in the 1800s, where pharmacists provided local traditional medicines to customers, which included extracts like morphine, quinine, and strychnine.^[95] Therapeutically important drugs like camptothecin (from *Camptotheca acuminata*, used in traditional Chinese medicine) and taxol (from the Pacific yew, *Taxus brevifolia*) were derived from medicinal plants.^{[96][34]} The Vinca alkaloids vincristine and vinblastine, used as anti-cancer drugs, were discovered in the 1950s from the Madagascar periwinkle, *Catharanthus roseus*.^[97]



The anticancer drug taxol was developed after screening of the Pacific yew, *Taxus brevifolia* (foliage and fruit shown).

Hundreds of compounds have been identified using ethnobotany, investigating plants used by indigenous peoples for possible medical applications.^[98] Some important phytochemicals, including curcumin, epigallocatechin gallate, genistein and resveratrol are pan-assay interference compounds, meaning that in vitro studies of their activity often provide unreliable data. As a result, phytochemicals have frequently proven unsuitable as lead compounds in drug discovery.^{[99][100]} In the United States over the period 1999 to 2012, despite several hundred applications for new drug status, only two botanical drug candidates had sufficient evidence of medicinal value to be approved by the Food and Drug Administration.^[2]

The pharmaceutical industry has remained interested in mining traditional uses of medicinal plants in its drug discovery efforts.^[34] Of the 1073 small-molecule drugs approved in the period 1981 to 2010, over half were either directly derived from or inspired by natural substances.^{[34][101]} Among cancer treatments, of 185 small-molecule drugs approved in the period from 1981 to 2019, 65% were derived from or inspired by natural substances.^[102]

Safety

Plant medicines can cause adverse effects and even death, whether by side-effects of their active substances, by adulteration or contamination, by overdose, or by inappropriate prescription. Many such effects are known, while others remain to be explored scientifically. There is no reason to presume that because a product comes from nature it must be safe: the existence of powerful natural poisons like atropine and nicotine shows this to be untrue. Further, the high standards applied to conventional medicines do not always apply to plant medicines, and dose can vary widely depending on the growth conditions of plants: older plants may be much more toxic than young ones, for instance.^{[104][105][106][107][108][109]}

Pharmacologically active plant extracts can interact with conventional drugs, both because they may provide an increased dose of similar compounds, and because some phytochemicals interfere with the body's systems that metabolise drugs in the liver including the cytochrome P450 system, making the drugs last longer in the body and have a more powerful cumulative effect.^[110] Plant medicines can be dangerous during pregnancy.^[111] Since plants may contain many different substances, plant extracts may have complex effects on the human body.^[5]



The Thornapple *Datura stramonium* has been used for asthma, because it contains the alkaloid atropine, but it is also a powerful and potentially fatal hallucinogen.^[103]

Quality, advertising, and labelling

Herbal medicine and dietary supplement products have been criticized as not having sufficient standards or scientific evidence to confirm their contents, safety, and presumed efficacy.^{[112][113][114][115]} A 2013 study found that one-third of herbal products sampled contained no trace of the herb listed on the label, and other products were adulterated with unlisted fillers including potential allergens.^{[116][117]}

Threats

Where medicinal plants are harvested from the wild rather than cultivated, they are subject to both general and specific threats. General threats include climate change and habitat loss to development and agriculture. A specific threat is over-collection to meet rising demand for medicines.^[118] A case in point was the pressure on wild populations of the Pacific yew soon after news of taxol's effectiveness became public.^[34] The threat from over-collection could be addressed by cultivation of some medicinal plants, or by a system of certification to make wild harvesting sustainable.^[118] A report in 2020 by the Royal Botanic Gardens, Kew identifies 723 medicinal plants as being at risk of extinction, caused partly by over-collection.^{[119][102]}

See also

- Ethnomedicine
- European Directive on Traditional Herbal Medicinal Products

Notes

- Farnsworth states that this figure was based on prescriptions from American community pharmacies between 1959 and 1980.^[38]
- Berberine is the main active component of an ancient Chinese herb *Coptis chinensis* French, which has been administered for what Yin and colleagues state is "diabetes" for thousands of years, although with no sound evidence of efficacy.^[54]

- c. Tobacco has "probably been responsible for more deaths than any other herb", but it was used as a medicine in the societies encountered by Columbus and was considered a panacea in Europe. It is no longer accepted as medicinal.^[55]

References

1. Lichterman, B. L. (2004). "Aspirin: The Story of a Wonder Drug" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC535471>). *British Medical Journal*. **329** (7479): 1408. doi:10.1136/bmj.329.7479.1408 (<https://doi.org/10.1136%2Fbmj.329.7479.1408>). PMC 535471 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC535471>).
2. Ahn, K. (2017). "The worldwide trend of using botanical drugs and strategies for developing global drugs" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5422022>). *BMB Reports*. **50** (3): 111–116. doi:10.5483/BMBRep.2017.50.3.221 (<https://doi.org/10.5483%2FBMBRep.2017.50.3.221>). PMC 5422022 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5422022>). PMID 27998396 (<https://pubmed.ncbi.nlm.nih.gov/27998396>).
3. "Medicinal and aromatic plants trade programme" (<https://web.archive.org/web/20180301133744/http://www.traffic.org/medicinal-plants/>). Traffic.org. Archived from the original (<http://www.traffic.org/medicinal-plants/>) on 1 March 2018. Retrieved 20 February 2017.
4. Collins, Minta (2000). *Medieval Herbals: The Illustrative Traditions* (<https://books.google.com/books?id=s7c2EaxDYjUC&pg=PA32>). University of Toronto Press. p. 32. ISBN 978-0-8020-8313-5.
5. Tapsell, L. C.; Hemphill, I.; Cobiac, L.; et al. (August 2006). "Health benefits of herbs and spices: the past, the present, the future" (<https://ro.uow.edu.au/cgi/viewcontent.cgi?article=2450&context=hbspapers>). *Med. J. Aust.* **185** (4 Suppl): S4–24. doi:10.5694/j.1326-5377.2006.tb00548.x (<https://doi.org/10.5694%2Fj.1326-5377.2006.tb00548.x>). PMID 17022438 (<https://pubmed.ncbi.nlm.nih.gov/17022438>). S2CID 9769230 (<https://api.semanticscholar.org/CorpusID:9769230>).
6. Billing, Jennifer; Sherman, P.W. (March 1998). "Antimicrobial functions of spices: why some like it hot" (<https://semanticscholar.org/paper/5dd3dda315f032dec522b5834b9d16c0e40b6e04>). *Q Rev Biol.* **73** (1): 3–49. doi:10.1086/420058 (<https://doi.org/10.1086%2F420058>). PMID 9586227 (<https://pubmed.ncbi.nlm.nih.gov/9586227>). S2CID 22420170 (<https://api.semanticscholar.org/CorpusID:22420170>).
7. Sherman, P.W.; Hash, G.A. (May 2001). "Why vegetable recipes are not very spicy". *Evol Hum Behav.* **22** (3): 147–163. doi:10.1016/S1090-5138(00)00068-4 (<https://doi.org/10.1016%2FS1090-5138%2800%2900068-4>). PMID 11384883 (<https://pubmed.ncbi.nlm.nih.gov/11384883>).
8. "Angiosperms: Division Magnoliophyta: General Features". *Encyclopædia Britannica* (volume 13, 15th edition). 1993. p. 609.
9. Stepp, John R. (June 2004). "The role of weeds as sources of pharmaceuticals". *Journal of Ethnopharmacology*. **92** (2–3): 163–166. doi:10.1016/j.jep.2004.03.002 (<https://doi.org/10.1016%2Fj.jep.2004.03.002>). PMID 15137997 (<https://pubmed.ncbi.nlm.nih.gov/15137997>).
10. Stepp, John R.; Moerman, Daniel E. (April 2001). "The importance of weeds in ethnopharmacology". *Journal of Ethnopharmacology*. **75** (1): 19–23. doi:10.1016/S0378-8741(00)00385-8 (<https://doi.org/10.1016%2FS0378-8741%2800%2900385-8>). PMID 11282438 (<https://pubmed.ncbi.nlm.nih.gov/11282438>).
11. Sumner, Judith (2000). *The Natural History of Medicinal Plants*. Timber Press. p. 16. ISBN 978-0-88192-483-1.
12. Solecki, Ralph S. (November 1975). "Shanidar IV, a Neanderthal Flower Burial in Northern Iraq". *Science*. **190** (4217): 880–881. Bibcode:1975Sci...190..880S (<https://ui.adsabs.harvard.edu/abs/1975Sci...190..880S>). doi:10.1126/science.190.4217.880 (<https://doi.org/10.1126%2Fscience.190.4217.880>). S2CID 71625677 (<https://api.semanticscholar.org/CorpusID:71625677>).

13. Capasso, L. (December 1998). "5300 years ago, the Ice Man used natural laxatives and antibiotics". *Lancet*. **352** (9143): 1864. doi:10.1016/S0140-6736(05)79939-6 (<https://doi.org/10.1016%2FS0140-6736%2805%2979939-6>). PMID 9851424 (<https://pubmed.ncbi.nlm.nih.gov/9851424>). S2CID 40027370 (<https://api.semanticscholar.org/CorpusID:40027370>).
14. Sumner, Judith (2000). *The Natural History of Medicinal Plants*. Timber Press. p. 17. ISBN 978-0-88192-483-1.
15. Aggarwal, B. B.; Sundaram, C.; Malani, N.; Ichikawa, H. (2007). *Curcumin: the Indian solid gold* (<https://archive.org/details/moleculartargets00bhar/page/1>). *Adv. Exp. Med. Biol.* Advances in Experimental Medicine and Biology. **595**. pp. 1–75 (<https://archive.org/details/moleculartargets00bhar/page/1>). doi:10.1007/978-0-387-46401-5_1 (https://doi.org/10.1007%2F978-0-387-46401-5_1). ISBN 978-0-387-46400-8. PMID 17569205 (<https://pubmed.ncbi.nlm.nih.gov/17569205>).
16. Girish Dwivedi, Shridhar Dwivedi (2007). *History of Medicine: Sushruta – the Clinician – Teacher par Excellence* (<https://web.archive.org/web/20081010045900/http://medind.nic.in/iae/t07/i4/iae/t07i4p243.pdf>) (PDF). National Informatics Centre. Archived from the original (<http://medind.nic.in/iae/t07/i4/iae/t07i4p243.pdf>) (PDF) on 10 October 2008. Retrieved 8 October 2008.
17. Sumner, Judith (2000). *The Natural History of Medicinal Plants*. Timber Press. p. 18. ISBN 978-0-88192-483-1.
18. Wu, Jing-Nuan (2005). *An Illustrated Chinese Materia Medica* (<https://books.google.com/books?id=H9Q8agQrJR0C&pg=PA6>). Oxford University Press. p. 6. ISBN 978-0-19-514017-0.
19. Grene, Marjorie (2004). *The philosophy of biology: an episodic history* (<https://books.google.com/books?id=TJSGJjh8Jo8C&pg=PA11>). Cambridge University Press. p. 11. ISBN 978-0-521-64380-1.
20. Arsdall, Anne V. (2002). *Medieval Herbal Remedies: The Old English Herbarium and Anglo-Saxon Medicine* (<https://books.google.com/books?id=ITxsagEmVHAC&pg=PA70>). Psychology Press. pp. 70–71. ISBN 978-0-415-93849-5.
21. Mills, Frank A. (2000). "Botany" (<https://books.google.com/books?id=GfC0TDkJJNgC&pg=PA179>). In Johnston, William M. (ed.). *Encyclopedia of Monasticism: M-Z*. Taylor & Francis. p. 179. ISBN 978-1-57958-090-2.
22. Ramos-e-Silva Marcia (1999). "Saint Hildegard Von Bingen (1098–1179) "The Light Of Her People And Of Her Time" ". *International Journal of Dermatology*. **38** (4): 315–320. doi:10.1046/j.1365-4362.1999.00617.x (<https://doi.org/10.1046%2Fj.1365-4362.1999.00617.x>). PMID 10321953 (<https://pubmed.ncbi.nlm.nih.gov/10321953>). S2CID 13404562 (<https://api.semanticscholar.org/CorpusID:13404562>).
23. Castleman, Michael (2001). *The New Healing Herbs* (<https://archive.org/details/newhealingherbs00mich/page/15>). Rodale. p. 15 (<https://archive.org/details/newhealingherbs00mich/page/15>). ISBN 978-1-57954-304-4.; Collins, Minta (2000). *Medieval Herbals: The Illustrative Traditions* (<https://books.google.com/books?id=s7c2EaxDYjUC&pg=PA115>). University of Toronto Press. p. 115. ISBN 978-0-8020-8313-5.; "Pharmaceutics and Alchemy" (https://www.nlm.nih.gov/exhibition/islamic_medical/islamic_11.html). US National Library of Medicine. Retrieved 26 January 2017.; Fahd, Toufic. "Botany and agriculture": 815. , in Rashed, Roshdi; Morelon, Régis (1996). *Encyclopedia of the History of Arabic Science: Astronomy-Theoretical and applied, v.2 Mathematics and the physical sciences; v.3 Technology, alchemy and life sciences* (<https://books.google.com/books?id=Fx7GtWAACAAJ>). Routledge. ISBN 978-0-415-02063-3.
24. Castleman, Michael (2001). *The New Healing Herbs* (<https://archive.org/details/newhealingherbs00mich/page/15>). Rodale. p. 15 (<https://archive.org/details/newhealingherbs00mich/page/15>). ISBN 978-1-57954-304-4.
25. Jacquart, Danielle (2008). "Islamic Pharmacology in the Middle Ages: Theories and Substances". *European Review*. **16** (2): 219–227 [223]. doi:10.1017/S1062798708000215 (<https://doi.org/10.1017%2FS1062798708000215>).

26. Kujundzić, E.; Masić, I. (1999). "[Al-Biruni--a universal scientist]". *Med. Arh.* (in Croatian). **53** (2): 117–120. PMID 10386051 (<https://pubmed.ncbi.nlm.nih.gov/10386051>).
27. Krek, M. (1979). "The Enigma of the First Arabic Book Printed from Movable Type". *Journal of Near Eastern Studies*. **38** (3): 203–212. doi:10.1086/372742 (<https://doi.org/10.1086%2F372742>). S2CID 162374182 (<https://api.semanticscholar.org/CorpusID:162374182>).
28. Brater, D. Craig & Daly, Walter J. (2000). "Clinical pharmacology in the Middle Ages: Principles that presage the 21st century". *Clinical Pharmacology & Therapeutics*. **67** (5): 447–450 [448–449]. doi:10.1067/mcp.2000.106465 (<https://doi.org/10.1067%2Fmcp.2000.106465>). PMID 10824622 (<https://pubmed.ncbi.nlm.nih.gov/10824622>). S2CID 45980791 (<https://api.semanticscholar.org/CorpusID:45980791>).
29. Singer, Charles (1923). "Herbals". *The Edinburgh Review*. **237**: 95–112.
30. Nunn, Nathan; Qian, Nancy (2010). "The Columbian Exchange: A History of Disease, Food, and Ideas". *Journal of Economic Perspectives*. **24** (2): 163–188. CiteSeerX 10.1.1.232.9242 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.232.9242>). doi:10.1257/jep.24.2.163 (<https://doi.org/10.1257%2Fjep.24.2.163>). JSTOR 25703506 (<https://www.jstor.org/stable/25703506>).
31. Heywood, Vernon H. (2012). "The role of New World biodiversity in the transformation of Mediterranean landscapes and culture" (<https://web.archive.org/web/20170227062205/http://147.163.105.223/bocconeae/24-069.pdf>) (PDF). *Bocconeae*. **24**: 69–93. Archived from the original (<http://147.163.105.223/bocconeae/24-069.pdf>) (PDF) on 2017-02-27. Retrieved 2017-02-26.
32. Gimmel Millie (2008). "Reading Medicine In The Codex De La Cruz Badiano". *Journal of the History of Ideas*. **69** (2): 169–192. doi:10.1353/jhi.2008.0017 (<https://doi.org/10.1353%2Fjhi.2008.0017>). PMID 19127831 (<https://pubmed.ncbi.nlm.nih.gov/19127831>). S2CID 46457797 (<https://api.semanticscholar.org/CorpusID:46457797>).
33. Petrovska 2012.
34. Atanasov, Atanas G.; Waltenberger, Birgit; Pferschy-Wenzig, Eva-Maria; Linder, Thomas; Wawrosch, Christoph; Uhrin, Pavel; Temml, Veronika; Wang, Limei; Schwaiger, Stefan; Heiss, Elke H.; Rollinger, Judith M.; Schuster, Daniela; Breuss, Johannes M.; Bochkov, Valery; Mihovilovic, Marko D.; Kopp, Brigitte; Bauer, Rudolf; Dirsch, Verena M.; Stuppner, Hermann (December 2015). "Discovery and resupply of pharmacologically active plant-derived natural products: A review" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4748402>). *Biotechnology Advances*. **33** (8): 1582–1614. doi:10.1016/j.biotechadv.2015.08.001 (<https://doi.org/10.1016%2Fj.biotechadv.2015.08.001>). PMC 4748402 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4748402>). PMID 26281720 (<https://pubmed.ncbi.nlm.nih.gov/26281720>).
35. Smith-Hall, C.; Larsen, H.O.; Pouliot, M. (2012). "People, plants and health: a conceptual framework for assessing changes in medicinal plant consumption" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3549945>). *J Ethnobiol Ethnomed*. **8**: 43. doi:10.1186/1746-4269-8-43 (<https://doi.org/10.1186%2F1746-4269-8-43>). PMC 3549945 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3549945>). PMID 23148504 (<https://pubmed.ncbi.nlm.nih.gov/23148504>).
36. Schippmann, Uwe; Leaman, Danna J.; Cunningham, A. B. (12 October 2002). "Impact of Cultivation and Gathering of Medicinal Plants on Biodiversity: Global Trends and Issues 2. Some Figures to start with ..." (<http://www.fao.org/docrep/005/AA010E/AA010e02.htm>) *Biodiversity and the Ecosystem Approach in Agriculture, Forestry and Fisheries. Satellite event on the occasion of the Ninth Regular Session of the Commission on Genetic Resources for Food and Agriculture. Rome, 12–13 October 2002. Inter-Departmental Working Group on Biological Diversity for Food and Agriculture. Rome.* Food and Agriculture Organization. Retrieved 25 September 2017.
37. "State of the World's Plants Report - 2016" (https://stateoftheworldsplants.com/2016/report/sotw_p_2016.pdf) (PDF). Royal Botanic Gardens, Kew. 2016. Retrieved 25 September 2017.

38. Farnsworth, Norman R.; Akerele, Olayiwola; Bingel, Audrey S.; Soejarto, Djada D.; Guo, Zhengang (1985). "Medicinal plants in therapy" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2536466>). *Bulletin of the World Health Organization*. **63** (6): 965–981. PMC 2536466 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2536466>). PMID 3879679 (<https://pubmed.ncbi.nlm.nih.gov/3879679>).
39. Tilburt, Jon C.; Kaptchuk, Ted J. (August 2008). "Herbal medicine research and global health: an ethical analysis" (<https://www.who.int/bulletin/volumes/86/8/07-042820/en/>). *Bulletin of the World Health Organization*. **86** (8): 577–656. doi:10.2471/BLT.07.042820 (<https://doi.org/10.2471/BLT.07.042820>). PMC 2649468 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2649468>). PMID 18797616 (<https://pubmed.ncbi.nlm.nih.gov/18797616>). Retrieved 22 September 2017.
40. Ekor, Martins (2013). "The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3887317>). *Frontiers in Pharmacology*. **4** (3): 202–4. doi:10.3389/fphar.2013.00177 (<https://doi.org/10.3389/fphar.2013.00177>). PMC 3887317 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3887317>). PMID 24454289 (<https://pubmed.ncbi.nlm.nih.gov/24454289>).
41. Singh, Amritpal (2016). *Regulatory and Pharmacological Basis of Ayurvedic Formulations* (<https://books.google.com/books?id=zgBPDAAAQBAJ&pg=PA4>). CRC Press. pp. 4–5. ISBN 978-1-4987-5096-7.
42. Cravotto, G.; Boffa, L.; Genzini, L.; Garella, D. (February 2010). "Phytotherapeutics: an evaluation of the potential of 1000 plants" (<https://doi.org/10.1111/j.1365-2710.2009.01096.x>). *Journal of Clinical Pharmacy and Therapeutics*. **35** (1): 11–48. doi:10.1111/j.1365-2710.2009.01096.x (<https://doi.org/10.1111/j.1365-2710.2009.01096.x>). PMID 20175810 (<https://pubmed.ncbi.nlm.nih.gov/20175810>). S2CID 29427595 (<https://api.semanticscholar.org/CorpusID:29427595>).
43. "Active Plant Ingredients Used for Medicinal Purposes" (<https://www.fs.fed.us/wildflowers/ethnobotany/medicinal/ingredients.shtml>). United States Department of Agriculture. Retrieved 18 February 2017. "Below are several examples of active plant ingredients that provide medicinal plant uses for humans."
44. Hayat, S. & Ahmad, A. (2007). *Salicylic Acid – A Plant Hormone* (<https://archive.org/details/salicylicacidpla0000unse>). Springer Science and Business Media. ISBN 978-1-4020-5183-8.
45. Bastida, Jaume; Lavilla, Rodolfo; Viladomat, Francesc Viladomat (2006). "Chemical and Biological Aspects of *Narcissus* Alkaloids". In Cordell, G. A. (ed.). *The Alkaloids: Chemistry and Biology*. **63**. pp. 87–179. doi:10.1016/S1099-4831(06)63003-4 ([https://doi.org/10.1016/S1099-4831\(06\)63003-4](https://doi.org/10.1016/S1099-4831(06)63003-4)). ISBN 978-0-12-469563-4. PMC 7118783 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7118783>). PMID 17133715 (<https://pubmed.ncbi.nlm.nih.gov/17133715>).
46. "Galantamine" (<https://www.drugs.com/mtm/galantamine.html>). Drugs.com. 2017. Retrieved 17 March 2018.
47. Birks, J. (2006). Birks, Jacqueline S (ed.). "Cholinesterase inhibitors for Alzheimer's disease". *The Cochrane Database of Systematic Reviews* (1): CD005593. doi:10.1002/14651858.CD005593 (<https://doi.org/10.1002/14651858.CD005593>). PMID 16437532 (<https://pubmed.ncbi.nlm.nih.gov/16437532>).
48. Soejarto, D. D. (1 March 2011). "Transcriptome Characterization, Sequencing, And Assembly Of Medicinal Plants Relevant To Human Health" (<http://apps.pharmacy.uic.edu/depts/pcrps/MedTranscriptomePlants/>). University of Illinois at Chicago. Retrieved 26 January 2017.
49. Meskin, Mark S. (2002). *Phytochemicals in Nutrition and Health* (<https://books.google.com/books?id=cJHsMALUDj0C&pg=PA123>). CRC Press. p. 123. ISBN 978-1-58716-083-7.
50. Springbob, Karen & Kutchan, Toni M. (2009). "Introduction to the different classes of natural products" (<https://books.google.com/books?id=Y8SpVXEng4QC&pg=PA3>). In Lanzotti, Virginia (ed.). *Plant-Derived Natural Products: Synthesis, Function, and Application*. Springer. p. 3. ISBN 978-0-387-85497-7.

51. Elumalai, A.; Eswariah, M. Chinna (2012). "Herbalism - A Review" (https://web.archive.org/web/20170217232814/http://www.phytotherapyjournal.com/File_Folder/96-105.pdf) (PDF). *International Journal of Phytotherapy*. **2** (2): 96–105. Archived from the original (http://www.phytotherapyjournal.com/File_Folder/96-105.pdf) (PDF) on 2017-02-17. Retrieved 2017-02-17.
52. Aniszewski, Tadeusz (2007). *Alkaloids – secrets of life*. Amsterdam: Elsevier. p. 182. ISBN 978-0-444-52736-3.
53. "Atropa Belladonna" (http://www.ema.europa.eu/docs/en_GB/document_library/Maximum_Residue_Limits_-_Report/2009/11/WC500010731.pdf) (PDF). The European Agency for the Evaluation of Medicinal Products. 1998. Retrieved 26 February 2017.
54. Yin, Jun; Xing, Huili; Ye, Jianping (May 2008). "Efficacy of Berberine in Patients with Type 2 Diabetes" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2410097>). *Metabolism*. **57** (5): 712–717. doi:10.1016/j.metabol.2008.01.013 (<https://doi.org/10.1016%2Fj.metabol.2008.01.013>). PMC 2410097 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2410097>). PMID 18442638 (<https://pubmed.ncbi.nlm.nih.gov/18442638>).
55. Charlton, Anne (June 2004). "Medicinal uses of tobacco in history" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1079499>). *Journal of the Royal Society of Medicine*. **97** (6): 292–296. doi:10.1258/jrsm.97.6.292 (<https://doi.org/10.1258%2Fjrsm.97.6.292>). PMC 1079499 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1079499>). PMID 15173337 (<https://pubmed.ncbi.nlm.nih.gov/15173337>).
56. Gremigni, P.; et al. (2003). "The interaction of phosphorus and potassium with seed alkaloid concentrations, yield and mineral content in narrow-leaved lupin (*Lupinus angustifolius* L.)". *Plant and Soil*. **253** (2): 413–427. doi:10.1023/A:1024828131581 (<https://doi.org/10.1023%2FA%3A1024828131581>). JSTOR 24121197 (<https://www.jstor.org/stable/24121197>). S2CID 25434984 (<https://api.semanticscholar.org/CorpusID:25434984>).
57. "Nicotinic acetylcholine receptors: Introduction" (<http://www.iuphar-db.org/DATABASE/FamilyIntroductionForward?familyId=76&familyType=IC>). *IUPHAR Database*. International Union of Basic and Clinical Pharmacology. Retrieved 26 February 2017.
58. Wang, Zhe; Ma, Pei; He, Chunnian; Peng, Yong; Xiao, Peigen (2013). "Evaluation of the content variation of anthraquinone glycosides in rhubarb by UPLC-PDA" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3854541>). *Chemistry Central Journal*. **7** (1): 43–56. doi:10.1186/1752-153X-7-170 (<https://doi.org/10.1186%2F1752-153X-7-170>). PMC 3854541 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3854541>). PMID 24160332 (<https://pubmed.ncbi.nlm.nih.gov/24160332>).
59. Chan, K.; Lin, T.X. (2009). *Treatments used in complementary and alternative medicine*. Side Effects of Drugs Annual. **31**. pp. 745–756. doi:10.1016/S0378-6080(09)03148-1 (<https://doi.org/10.1016%2FS0378-6080%2809%2903148-1>). ISBN 978-0-444-53294-7.
60. Hietala, P.; Marvola, M.; Parviainen, T.; Lainonen, H. (August 1987). "Laxative potency and acute toxicity of some anthraquinone derivatives, senna extracts and fractions of senna extracts". *Pharmacology & Toxicology*. **61** (2): 153–6. doi:10.1111/j.1600-0773.1987.tb01794.x (<https://doi.org/10.1111%2Fj.1600-0773.1987.tb01794.x>). PMID 3671329 (<https://pubmed.ncbi.nlm.nih.gov/3671329>).
61. Akolkar, Praful (2012-12-27). "Pharmacognosy of Rhubarb" (<http://pharmaxchange.info/press/2012/12/pharmacognosy-of-rhubarb/>). *PharmaXChange.info*.
62. "Digitalis purpurea. Cardiac Glycoside" (<http://botany.csd.tamu.edu/FLORA/Wilson/481/medbot/bot2.htm>). Texas A&M University. Retrieved 26 February 2017. "The man credited with the introduction of digitalis into the practice of medicine was William Withering."
63. Da Silva, Cecilia; et al. (2013). "The High Polyphenol Content of Grapevine Cultivar Tannat Berries Is Conferred Primarily by Genes That Are Not Shared with the Reference Genome" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3903987>). *The Plant Cell*. **25** (12): 4777–4788. doi:10.1105/tpc.113.118810 (<https://doi.org/10.1105%2Ftpc.113.118810>). JSTOR 43190600 (<https://www.jstor.org/stable/43190600>). PMC 3903987 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3903987>). PMID 24319081 (<https://pubmed.ncbi.nlm.nih.gov/24319081>).

64. Muller-Schwarze, Dietland (2006). *Chemical Ecology of Vertebrates*. Cambridge University Press. p. 287. ISBN 978-0-521-36377-8.
65. Lee, Y. S.; Park J. S.; Cho S. D.; Son J. K.; Cherdshewasart W.; Kang K. S. (Dec 2002). "Requirement of metabolic activation for estrogenic activity of *Pueraria mirifica*" (<http://www.koreascience.or.kr/article/JAKO200210103477199.page>). *Journal of Veterinary Science*. **3** (4): 273–277. CiteSeerX 10.1.1.617.1507 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.617.1507>). doi:10.4142/jvs.2002.3.4.273 (<https://doi.org/10.4142%2Fjvs.2002.3.4.273>). PMID 12819377 (<https://pubmed.ncbi.nlm.nih.gov/12819377>).
66. Delmonte, P.; Rader, J. I. (2006). "Analysis of isoflavones in foods and dietary supplements" (<https://doi.org/10.1093%2Fjaoac%2F89.4.1138>). *Journal of AOAC International*. **89** (4): 1138–46. doi:10.1093/jaoac/89.4.1138 (<https://doi.org/10.1093%2Fjaoac%2F89.4.1138>). PMID 16915857 (<https://pubmed.ncbi.nlm.nih.gov/16915857>).
67. Brown, D.E.; Walton, N. J. (1999). *Chemicals from Plants: Perspectives on Plant Secondary Products*. World Scientific Publishing. pp. 21, 141. ISBN 978-981-02-2773-9.
68. Albert-Puleo, M. (Dec 1980). "Fennel and anise as estrogenic agents". *Journal of Ethnopharmacology*. **2** (4): 337–44. doi:10.1016/S0378-8741(80)81015-4 ([https://doi.org/10.1016/S0378-8741\(80\)81015-4](https://doi.org/10.1016%2FS0378-8741%2880%2981015-4)). PMID 6999244 (<https://pubmed.ncbi.nlm.nih.gov/6999244>).
69. European Food Safety Authority (2010). "Scientific Opinion on the substantiation of health claims related to various food(s)/food constituent(s) and protection of cells from premature aging, antioxidant activity, antioxidant content and antioxidant properties, and protection of DNA, proteins and lipids from oxidative damage pursuant to Article 13(1) of Regulation (EC) No 1924/2006" (<https://doi.org/10.2903%2Fj.efsa.2010.1489>). *EFSA Journal*. **8** (2): 1489. doi:10.2903/j.efsa.2010.1489 (<https://doi.org/10.2903%2Fj.efsa.2010.1489>).
70. Jindal, K. K.; Sharma, R. C. (2004). *Recent trends in horticulture in the Himalayas* (<https://books.google.com/books?id=LlogqveEFVgC>). Indus Publishing. ISBN 978-81-7387-162-7.
71. Turner, J. V.; Agatonovic-Kustrin, S.; Glass, B. D. (Aug 2007). "Molecular aspects of phytoestrogen selective binding at estrogen receptors". *Journal of Pharmaceutical Sciences*. **96** (8): 1879–85. doi:10.1002/jps.20987 (<https://doi.org/10.1002%2Fjps.20987>). PMID 17518366 (<https://pubmed.ncbi.nlm.nih.gov/17518366>).
72. Wiart, Christopher (2014). *Terpenes. Lead Compounds from Medicinal Plants for the Treatment of Neurodegenerative Diseases*. pp. 189–284.
73. Tchen, T. T. (1965). "Reviewed Work: *The Biosynthesis of Steroids, Terpenes & Acetogenins*". *American Scientist*. **53** (4): 499A–500A. JSTOR 27836252 (<https://www.jstor.org/stable/27836252>).
74. Singsaas, Eric L. (2000). "Terpenes and the Thermotolerance of Photosynthesis" (<https://doi.org/10.1046%2Fj.1469-8137.2000.00626.x>). *New Phytologist*. **146** (1): 1–2. doi:10.1046/j.1469-8137.2000.00626.x (<https://doi.org/10.1046%2Fj.1469-8137.2000.00626.x>). JSTOR 2588737 (<https://www.jstor.org/stable/2588737>).
75. "Thymol (CID=6989)" (<https://pubchem.ncbi.nlm.nih.gov/compound/thymol>). NIH. Retrieved 26 February 2017. "THYMOL is a phenol obtained from thyme oil or other volatile oils used as a stabilizer in pharmaceutical preparations, and as an antiseptic (antibacterial or antifungal) agent. It was formerly used as a vermifuge."
76. "WHO Guidelines on Good Agricultural and Collection Practices (GACP) for Medicinal Plants" (<http://apps.who.int/medicinedocs/en/d/Js4928e/4.3.html>). World Health Organization. 2003. Retrieved 26 February 2017.
77. Carrubba, A.; Scalenghe, R. (2012). "Scent of Mare Nostrum — Medicinal and Aromatic Plants (MAPs) in Mediterranean soils". *Journal of the Science of Food and Agriculture*. **92** (6): 1150–1170. doi:10.1002/jsfa.5630 (<https://doi.org/10.1002%2Fjsfa.5630>). PMID 22419102 (<https://pubmed.ncbi.nlm.nih.gov/22419102>).

78. Yang, Yifan (2010). "Theories and concepts in the composition of Chinese herbal formulas" (<https://www.sciencedirect.com/science/article/pii/B9780702031328000062>). *Chinese Herbal Formulas*. Elsevier Ltd.: 1–34. doi:10.1016/B978-0-7020-3132-8.00006-2 (<https://doi.org/10.1016%2FB978-0-7020-3132-8.00006-2>). ISBN 9780702031328. Retrieved 18 April 2020.
79. Dharmananda, Subhuti (May 1997). "The Methods of Preparation of Herb Formulas: Decoctions, Dried Decoctions, Powders, Pills, Tablets, and Tinctures" (<http://www.itmonline.org/arts/methprep.htm>). Institute of Traditional Medicine, Portland, Oregon.
80. Mount, Toni (20 April 2015). "9 weird medieval medicines" (<http://www.historyextra.com/article/medieval/9-weird-medieval-medicines>). British Broadcasting Corporation.
81. Pezzuto, John M. (January 1997). "Plant-derived anticancer agents". *Biochemical Pharmacology*. **53** (2): 121–133. doi:10.1016/S0006-2952(96)00654-5 (<https://doi.org/10.1016%2FS0006-2952%2896%2900654-5>). PMID 9037244 (<https://pubmed.ncbi.nlm.nih.gov/9037244>).
82. "Traditional Medicine. Fact Sheet No. 134" (<https://www.webcitation.org/5Zeyw2hfS?url=https://www.who.int/mediacentre/factsheets/fs134/en/>). World Health Organization. May 2003. Archived from the original (<https://www.who.int/mediacentre/factsheets/fs134/en/>) on 28 July 2008. Retrieved 26 February 2017.
83. Chan, Margaret (19 August 2015). "WHO Director-General addresses traditional medicine forum" (<https://www.who.int/dg/speeches/2015/traditional-medicine/en/>). WHO.
84. "Traditional Chinese Medicine: In Depth (D428)" (<https://nccih.nih.gov/health/whatiscam/chinesemed.htm>). NIH. April 2009. Retrieved 26 February 2017.
85. Giovannini, Peter. "Managing diabetes with medicinal plants" (<https://www.kew.org/blogs/kew-science/managing-diabetes-medicinal-plants-0>). Kew Gardens. Retrieved 3 October 2017.
86. Giovannini, Peter; Howes, Melanie-Jayne R.; Edwards, Sarah E. (2016). "Medicinal plants used in the traditional management of diabetes and its sequelae in Central America: A review" ([https://kclpure.kcl.ac.uk/portal/en/publications/medicinal-plants-used-in-the-traditional-management-of-diabetes-and-its-sequelae-in-central-america-a-review\(69685ce6-6956-4937-bf12-ef28f0c3144c\).html](https://kclpure.kcl.ac.uk/portal/en/publications/medicinal-plants-used-in-the-traditional-management-of-diabetes-and-its-sequelae-in-central-america-a-review(69685ce6-6956-4937-bf12-ef28f0c3144c).html)). *Journal of Ethnopharmacology*. **184**: 58–71. doi:10.1016/j.jep.2016.02.034 (<https://doi.org/10.1016%2Fj.jep.2016.02.034>). PMID 26924564 (<https://pubmed.ncbi.nlm.nih.gov/26924564>).
87. Milliken, William (2015). "Medicinal knowledge in the Amazon" (<https://www.kew.org/blogs/kew-science/medicinal-knowledge-amazon>). Kew Gardens.
88. Yanomami, M. I.; Yanomami, E.; Albert, B.; Milliken, W; Coelho, V. (2014). *Hwëri mamotima thëpë ã oni. Manual dos remédios tradicionais Yanomami [Manual of Traditional Yanomami Medicines]*. São Paulo: Hutukara/Instituto Socioambiental.
89. "Scoring drugs. A new study suggests alcohol is more harmful than heroin or crack" (https://www.economist.com/blogs/dailychart/2010/11/drugs_cause_most_harm). *The Economist*. 2 November 2010. Retrieved 26 February 2017. ""Drug harms in the UK: a multi-criteria decision analysis", by David Nutt, Leslie King and Lawrence Phillips, on behalf of the Independent Scientific Committee on Drugs. *The Lancet*."
90. "Herbal medicine" (<https://www.cancerresearchuk.org/about-cancer/cancer-in-general/treatment/complementary-alternative-therapies/individual-therapies/herbal-medicine>). Cancer Research UK. Retrieved 7 July 2019. "There is no reliable evidence from human studies that herbal remedies can treat, prevent or cure any type of cancer. Some clinical trials seem to show that certain Chinese herbs may help people to live longer, might reduce side effects, and help to prevent cancer from coming back. This is especially when combined with conventional treatment."

91. Saslis-Lagoudakis, C. H.; Savolainen, V.; Williamson, E. M.; Forest, F.; Wagstaff, S. J.; Baral, S. R.; Watson, M. F.; Pendry, C. A.; Hawkins, J. A. (2012). "Phylogenies reveal predictive power of traditional medicine in bioprospecting" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3465383>). *Proceedings of the National Academy of Sciences*. **109** (39): 15835–40. Bibcode:2012PNAS..10915835S (<https://ui.adsabs.harvard.edu/abs/2012PNAS..10915835S>). doi:10.1073/pnas.1202242109 (<https://doi.org/10.1073/pnas.1202242109>). PMC 3465383 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3465383>). PMID 22984175 (<https://pubmed.ncbi.nlm.nih.gov/22984175>).
92. "International Regulatory Cooperation for Herbal Medicines (IRCH)" (<http://who.int/medicines/areas/traditional/irch/en/>). World Health Organization. Retrieved 2 October 2017.
93. Kala, Chandra Prakash; Sajwan, Bikram Singh (2007). "Revitalizing Indian systems of herbal medicine by the National Medicinal Plants Board through institutional networking and capacity building". *Current Science*. **93** (6): 797–806. JSTOR 24099124 (<https://www.jstor.org/stable/24099124>).
94. World Health Organization (2013). *WHO Traditional Medicine Strategy 2014-2023* (http://apps.who.int/iris/bitstream/10665/92455/1/9789241506090_eng.pdf?ua=1) (PDF). World Health Organization. ISBN 978-92-4-150609-0.
95. "Emergence of Pharmaceutical Science and Industry: 1870-1930" (<http://pubs.acs.org/cen/covestory/83/8325/8325emergence.html>). *Chemical & Engineering News*. **83** (25). 20 June 2005.
96. Heinrich, M.; Bremner, P. (March 2006). "Ethnobotany and ethnopharmacy--their role for anti-cancer drug development". *Current Drug Targets*. **7** (3): 239–245. doi:10.2174/138945006776054988 (<https://doi.org/10.2174/138945006776054988>). PMID 16515525 (<https://pubmed.ncbi.nlm.nih.gov/16515525>).
97. Moudi, Maryam; Go, Rusea; Yien, Christina Yong Seok; Nazre, Mohd. (November 2013). "Vinca Alkaloids" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883245>). *International Journal of Preventive Medicine*. **4** (11): 1231–1235. PMC 3883245 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883245>). PMID 24404355 (<https://pubmed.ncbi.nlm.nih.gov/24404355>).
98. Fabricant, D. S.; Farnsworth, N. R. (March 2001). "The value of plants used in traditional medicine for drug discovery" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1240543>). *Environ. Health Perspect.* **109** (Suppl 1): 69–75. doi:10.1289/ehp.01109s169 (<https://doi.org/10.1289/ehp.01109s169>). PMC 1240543 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1240543>). PMID 11250806 (<https://pubmed.ncbi.nlm.nih.gov/11250806>).
99. Baell, Jonathan; Walters, Michael A. (24 September 2014). "Chemistry: Chemical con artists foil drug discovery" (<https://doi.org/10.1038/2F513481a>). *Nature*. **513** (7519): 481–483. Bibcode:2014Natur.513..481B (<https://ui.adsabs.harvard.edu/abs/2014Natur.513..481B>). doi:10.1038/513481a (<https://doi.org/10.1038/2F513481a>). PMID 25254460 (<https://pubmed.ncbi.nlm.nih.gov/25254460>).
00. Dahlin, Jayme L; Walters, Michael A (July 2014). "The essential roles of chemistry in high-throughput screening triage" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4465542>). *Future Medicinal Chemistry*. **6** (11): 1265–90. doi:10.4155/fmc.14.60 (<https://doi.org/10.4155/fmc.14.60>). PMC 4465542 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4465542>). PMID 25163000 (<https://pubmed.ncbi.nlm.nih.gov/25163000>).
01. Newman, David J.; Cragg, Gordon M. (8 February 2012). "Natural Products As Sources of New Drugs over the 30 Years from 1981 to 2010" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3721181>). *Journal of Natural Products*. **75** (3): 311–35. doi:10.1021/np200906s (<https://doi.org/10.1021/np200906s>). PMC 3721181 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3721181>). PMID 22316239 (<https://pubmed.ncbi.nlm.nih.gov/22316239>).
02. "State of the World's Plants and Fungi 2020" (<https://www.kew.org/sites/default/files/2020-09/Kew%20State%20of%20the%20Worlds%20Plants%20and%20Fungi.pdf>) (PDF). Royal Botanic Gardens, Kew. 2020. Retrieved 30 September 2020.

03. Freye, Enno (2010). "Toxicity of Datura Stramonium". *Pharmacology and Abuse of Cocaine, Amphetamines, Ecstasy and Related Designer Drugs*. Springer. pp. 217–218. doi:10.1007/978-90-481-2448-0_34 (https://doi.org/10.1007%2F978-90-481-2448-0_34). ISBN 978-90-481-2447-3.
04. Ernst, E. (1998). "Harmless Herbs? A Review of the Recent Literature" (<http://toxicology.usu.edu/endnote/Harmless-Herbs.pdf>) (PDF). *The American Journal of Medicine*. **104** (2): 170–178. doi:10.1016/S0002-9343(97)00397-5 (<https://doi.org/10.1016%2FS0002-9343%2897%2900397-5>). PMID 9528737 (<https://pubmed.ncbi.nlm.nih.gov/9528737>).
05. Talalay, P. (2001). "The importance of using scientific principles in the development of medicinal agents from plants" (<https://doi.org/10.1097%2F00001888-200103000-00010>). *Academic Medicine*. **76** (3): 238–47. doi:10.1097/00001888-200103000-00010 (<https://doi.org/10.1097%2F00001888-200103000-00010>). PMID 11242573 (<https://pubmed.ncbi.nlm.nih.gov/11242573>).
06. Elvin-Lewis, M. (2001). "Should we be concerned about herbal remedies". *Journal of Ethnopharmacology*. **75** (2–3): 141–164. doi:10.1016/S0378-8741(00)00394-9 (<https://doi.org/10.1016%2FS0378-8741%2800%2900394-9>). PMID 11297844 (<https://pubmed.ncbi.nlm.nih.gov/11297844>).
07. Vickers, A. J. (2007). "Which botanicals or other unconventional anticancer agents should we take to clinical trial?" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2590766>). *J Soc Integr Oncol*. **5** (3): 125–9. doi:10.2310/7200.2007.011 (<https://doi.org/10.2310%2F7200.2007.011>). PMC 2590766 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2590766>). PMID 17761132 (<https://pubmed.ncbi.nlm.nih.gov/17761132>).
08. Ernst, E. (2007). "Herbal medicines: balancing benefits and risks". *Dietary Supplements and Health. Novartis Found. Symp. Novartis Foundation Symposia*. **282**. pp. 154–67, discussion 167–72, 212–8. doi:10.1002/9780470319444.ch11 (<https://doi.org/10.1002%2F9780470319444.ch11>). ISBN 978-0-470-31944-4. PMID 17913230 (<https://pubmed.ncbi.nlm.nih.gov/17913230>).
09. Pinn, G. (November 2001). "Adverse effects associated with herbal medicine". *Aust Fam Physician*. **30** (11): 1070–5. PMID 11759460 (<https://pubmed.ncbi.nlm.nih.gov/11759460>).
10. Nekvindová, J.; Anzenbacher, P. (July 2007). "Interactions of food and dietary supplements with drug metabolising cytochrome P450 enzymes". *Ceska Slov Farm*. **56** (4): 165–73. PMID 17969314 (<https://pubmed.ncbi.nlm.nih.gov/17969314>).
11. Born, D.; Barron, ML (May 2005). "Herb use in pregnancy: what nurses should know". *MCN Am J Matern Child Nurs*. **30** (3): 201–6. doi:10.1097/00005721-200505000-00009 (<https://doi.org/10.1097%2F00005721-200505000-00009>). PMID 15867682 (<https://pubmed.ncbi.nlm.nih.gov/15867682>). S2CID 35882289 (<https://api.semanticscholar.org/CorpusID:35882289>).
12. Barrett, Stephen (23 November 2013). "The herbal minefield" (<https://www.quackwatch.com/01/QuackeryRelatedTopics/herbs.html>). Quackwatch.
13. Zhang, J.; Wider, B.; Shang, H.; Li, X.; Ernst, E. (2012). "Quality of herbal medicines: Challenges and solutions". *Complementary Therapies in Medicine*. **20** (1–2): 100–106. doi:10.1016/j.ctim.2011.09.004 (<https://doi.org/10.1016%2Fj.ctim.2011.09.004>). PMID 22305255 (<https://pubmed.ncbi.nlm.nih.gov/22305255>).
14. Morris, C. A.; Avorn, J. (2003). "Internet marketing of herbal products". *JAMA*. **290** (11): 1505–9. doi:10.1001/jama.290.11.1505 (<https://doi.org/10.1001%2Fjama.290.11.1505>). PMID 13129992 (<https://pubmed.ncbi.nlm.nih.gov/13129992>).
15. Coghlan, M. L.; Haile, J.; Houston, J.; Murray, D. C.; White, N. E.; Moolhuijzen, P; Bellgard, M. I.; Bunce, M (2012). "Deep Sequencing of Plant and Animal DNA Contained within Traditional Chinese Medicines Reveals Legality Issues and Health Safety Concerns" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3325194>). *PLOS Genetics*. **8** (4): e1002657. doi:10.1371/journal.pgen.1002657 (<https://doi.org/10.1371%2Fjournal.pgen.1002657>). PMC 3325194 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3325194>). PMID 22511890 (<https://pubmed.ncbi.nlm.nih.gov/22511890>).

16. Newmaster, S. G.; Grguric, M.; Shanmughanandhan, D.; Ramalingam, S.; Ragupathy, S. (2013). "DNA barcoding detects contamination and substitution in North American herbal products" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3851815>). *BMC Medicine*. **11**: 222. doi:10.1186/1741-7015-11-222 (<https://doi.org/10.1186%2F1741-7015-11-222>). PMC 3851815 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3851815>). PMID 24120035 (<https://pubmed.ncbi.nlm.nih.gov/24120035>).
17. O'Connor, Anahad (5 November 2013). "Herbal Supplements Are Often Not What They Seem" (<https://www.nytimes.com/2013/11/05/science/herbal-supplements-are-often-not-what-they-see.html>). *The New York Times*.
18. Kling, Jim (2016). "Protecting medicine's wild pharmacy". *Nature Plants*. **2** (5): 16064. doi:10.1038/nplants.2016.64 (<https://doi.org/10.1038%2Fnplants.2016.64>). PMID 27243657 (<https://pubmed.ncbi.nlm.nih.gov/27243657>). S2CID 7246069 (<https://api.semanticscholar.org/CorpusID:7246069>).
19. Briggs, Helen. "Two-fifths of plants at risk of extinction, says report" (<https://www.bbc.co.uk/news/science-environment-54344309>). *BBC*. Retrieved 30 September 2020.

Further reading

- Aronson, Jeffrey K. (2008). *Meyler's Side Effects of Herbal Medicines* (https://books.google.com/books?id=_hkSTiEsKvkC). Elsevier. ISBN 978-0-08-093290-3.
- Braun, Lesley; Cohen, Marc (2007). *Herbs and Natural Supplements: An Evidence-Based Guide*. Elsevier. ISBN 978-0-7295-3796-4.
- Crellin, J. K.; et al. (199). *Herbal Medicine Past and Present: A reference guide to medicinal plants* (<https://books.google.com/books?id=0JaqB07uTx4C>). Duke University Press. ISBN 978-0-8223-1019-8.
- Lewis, Walter H. (2003). *Medical Botany: Plants Affecting Human Health* (<https://books.google.com/books?id=ipQmSriMF9sC>). John Wiley & Sons. ISBN 978-0-471-62882-8.
- Newall, Carol A.; et al. (1996). *Herbal medicines: a guide for health-care professionals* (https://books.google.com/books?id=n_psAAAAMAAJ). Pharmaceutical Press. ISBN 978-0-85369-289-8.
- Petrovska, Biljana Bauer (2012). "Historical review of medicinal plants' usage" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3358962>). *Pharmacognosy Reviews*. **6** (11): 1–5. doi:10.4103/0973-7847.95849 (<https://doi.org/10.4103%2F0973-7847.95849>). PMC 3358962 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3358962>). PMID 22654398 (<https://pubmed.ncbi.nlm.nih.gov/22654398>).

Retrieved from "https://en.wikipedia.org/w/index.php?title=Medicinal_plants&oldid=1019210153"

This page was last edited on 22 April 2021, at 02:48 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.