


Positional notation

Positional notation (or **place-value notation**, or **positional numeral system**) usually denotes the extension to any **base** of the **Hindu–Arabic numeral system** (or **decimal system**). More generally, a positional system is a numeral system in which the contribution of a digit to the value of a number is the value of the digit multiplied by a factor determined by the position of the digit. In early **numeral systems**, such as **Roman numerals**, a digit has only one value: I means one, X means ten and C a hundred (however, the value may be negated if placed before another digit). In modern positional systems, such as the **decimal system**, the position of the digit means that its value must be multiplied by some value: in 555, the three identical symbols represent five hundreds, five tens, and five units, respectively, due to their different positions in the digit string.




Glossary of terms used in the positional numeral systems

The [Babylonian numeral system](#), base 60, was the first positional system to be developed, and its influence is present today in the way time and angles are counted in tallies related to 60, such as 60 minutes in an hour and 360 degrees in a circle. Today, the Hindu–Arabic numeral system ([base ten](#)) is the most commonly used system globally. However, the [binary numeral system](#) (base two) is used in almost all [computers](#) and [electronic devices](#) because it is easier to implement efficiently in [electronic circuits](#).

Systems with negative base, [complex](#) base or negative digits have been described. Most of them do not require a minus sign for designating negative numbers.

The use of a [radix point](#) (decimal point in base ten), extends to include [fractions](#) and allows representing any [real number](#) with arbitrary accuracy. With positional notation, [arithmetical computations](#) are much simpler than with any older numeral system; this led to the rapid spread of the notation when it was introduced in western Europe.

History



Suanpan (the number represented in the picture is 6,302,715,408)

Today, the base-10 ([decimal](#)) system, which is presumably motivated by counting with the ten [fingers](#), is ubiquitous. Other bases have been used in the past, and some continue to be used today. For example, the [Babylonian numeral system](#), credited as the first positional numeral system, was [base-60](#). However, it lacked a real [zero](#). Initially inferred only from context, later, by about 700 BC, zero came to be indicated by a "space" or a "punctuation symbol" (such as two slanted wedges) between numerals.^[1] It was a [placeholder](#) rather than a true zero because it was not used alone or at the end of a number. Numbers like 2 and 120 (2×60) looked the same because the larger number lacked a final placeholder. Only context could differentiate them.

The polymath [Archimedes](#) (ca. 287–212 BC) invented a decimal positional system in his [Sand Reckoner](#) which was based on 10^8 ^[2] and later led the German mathematician [Carl Friedrich Gauss](#) to lament what heights science would have already reached in his days if Archimedes had fully realized the potential of his ingenious discovery.^[3]

Before positional notation became standard, simple additive systems ([sign-value notation](#)) such as [Roman numerals](#) were used, and accountants in ancient Rome and during the Middle Ages used the [abacus](#) or stone counters to do arithmetic.^[4]

1	2	3	4	5	6	7	8	9
					⊥	⊥	⊥	⊥
—	=	≡	≡	≡	⊥	⊥	⊥	⊥

*Chinese [rod numerals](#); Upper row vertical form
Lower row horizontal form*

[Counting rods](#) and most abacuses have been used to represent numbers in a positional numeral system. With counting rods or [abacus](#) to perform arithmetic operations, the writing of the starting, intermediate and final values of a calculation could easily be done with a simple additive system in each position or column. This approach required no memorization of tables (as does positional notation) and could produce practical results quickly.

The oldest extant positional notation system is that of Chinese [rod numerals](#), used from at least the early 8th century. It isn't clear whether this system was introduced from India or whether it was developed locally. [Indian numerals](#) originate with the [Brahmi numerals](#) of about the 3rd century BC, which symbols were, at the time, not used positionally. Medieval Indian numerals are positional, as are the derived [Arabic numerals](#), recorded from the 10th century.

After the [French Revolution](#) (1789–1799), the new French government promoted the extension of the decimal system.^[5] Some of those pro-decimal efforts—such as [decimal time](#) and the [decimal calendar](#)—were unsuccessful. Other French pro-decimal efforts—currency [decimalisation](#) and the [metrication](#) of weights and measures—spread widely out of France to almost the whole world.

History of positional fractions

J. Lennart Berggren notes that positional decimal fractions were used for the first time by Arab mathematician [Abu'l-Hasan al-Uqlidisi](#) as early as the 10th century.^[6] The Jewish mathematician [Immanuel Bonfils](#) used decimal fractions around 1350, but did not develop any notation to represent them.^[7] The Persian mathematician [Jamshīd al-Kāshī](#) made the same discovery of decimal fractions in the 15th century.^[6] [Al Khwarizmi](#) introduced fractions to Islamic countries in the early 9th century; his fraction presentation was similar to the traditional Chinese mathematical fractions from [Sunzi Suanjing](#).^[8] This form of fraction with numerator on top and denominator at bottom without a horizontal bar was also used by 10th century [Abu'l-Hasan al-Uqlidisi](#) and 15th century [Jamshīd al-Kāshī](#)'s work "Arithmetic Key".^{[8][9]}

The adoption of the decimal representation of numbers less than

Number: 184.54290

Simon Stevin's notation: 184①5①4②2③9④0

one, a [fraction](#), is often credited to [Simon Stevin](#) through his textbook [De Thiende](#),^[10] but both Stevin and [E. J. Dijksterhuis](#) indicate that [Regiomontanus](#) contributed to the European adoption of general [decimals](#).^[11]

European mathematicians, when taking over from the Hindus, *via* the Arabs, the idea of positional value for integers, neglected to extend this idea to fractions. For some centuries they confined themselves to using common and [sexagesimal](#) fractions... This half-heartedness has never been completely overcome, and sexagesimal fractions still form the basis of our trigonometry, astronomy and measurement of time. ¶ ... Mathematicians sought to avoid fractions by taking the radius R equal to a number of units of length of the form 10^n and then assuming for n so great an integral value that all occurring quantities could be expressed with sufficient accuracy by integers. ¶ The first to apply this method was the German astronomer Regiomontanus. To the extent that he expressed goniometrical line-segments in a unit $R/10^n$, Regiomontanus may be called an anticipator of the doctrine of decimal positional fractions.^{[11]:17,18}

In the estimation of Dijksterhuis, "after the publication of [De Thiende](#) only a small advance was required to establish the complete system of decimal positional fractions, and this step was taken promptly by a number of writers ... next to Stevin the most important figure in this development was Regiomontanus." Dijksterhuis noted that [Stevin] "gives full credit to

Regiomontanus for his prior contribution, saying that the trigonometric tables of the German astronomer actually contain the whole theory of 'numbers of the tenth progress'.^{[11]:19}

Issues

A key argument against the positional system was its susceptibility to easy [fraud](#) by simply putting a number at the beginning or end of a quantity, thereby changing (e.g.) 100 into 5100, or 100 into 1000. Modern [cheques](#) require a natural language spelling of an amount, as well as the decimal amount itself, to prevent such fraud. For the same reason the Chinese also use natural language numerals, for example 100 is written as 壹佰, which can never be forged into 壹仟 (1000) or 伍仟壹佰(5100).

Many of the advantages claimed for the metric system could be realized by any consistent positional notation. [Dozenal advocates](#) say duodecimal has several advantages over decimal, although the [switching cost](#) appears to be high.

Mathematics

Base of the numeral system

In [mathematical numeral systems](#) the radix r is usually the number of unique [digits](#), including zero, that a positional numeral system uses to represent numbers. In the interesting cases the radix is the [absolute value](#) $r = |b|$ of the base b , which may also be negative. For example, for the decimal system the radix (and base) is ten, because it uses the ten digits from 0 through 9. When a number "hits" 9, the next number will not be another different symbol, but a "1" followed by a "0". In binary, the radix is two, since after it hits "1", instead of "2" or another written symbol, it jumps straight to "10", followed by "11" and "100".

The highest symbol of a positional numeral system usually has the value one less than the value of the radix of that numeral system. The standard positional numeral systems differ from one another only in the base they use.

The radix is an integer that is greater than 1, since a radix of zero would not have any digits, and a radix of 1 would only have the zero digit. Negative bases are rarely used. In a system with more than $|b|$ unique digits, numbers may have many different possible representations.

It is important that the radix is finite, from which follows that the number of digits is quite low. Otherwise, the length of a numeral would not necessarily be [logarithmic](#) in its size.

(In certain [non-standard positional numeral systems](#), including [bijective numeration](#), the definition of the base or the allowed digits deviates from the above.)

In standard base-ten (decimal) positional notation, there are ten [decimal digits](#) and the number

$$5305_{\text{dec}} = (5 \times 10^3) + (3 \times 10^2) + (0 \times 10^1) + (5 \times 10^0).$$

In standard base-sixteen ([hexadecimal](#)), there are the sixteen hexadecimal digits (0–9 and A–F) and the number

$$14\text{B}9_{\text{hex}} = (1 \times 16^3) + (4 \times 16^2) + (\text{B} \times 16^1) + (9 \times 16^0) \quad (= 5305_{\text{dec}}),$$

where B represents the number eleven as a single symbol.

In general, in base- b , there are b digits $\{d_1, d_2, \dots, d_b\} =: D$ and the number

$$(a_3 a_2 a_1 a_0)_b = (a_3 \times b^3) + (a_2 \times b^2) + (a_1 \times b^1) + (a_0 \times b^0)$$

has $\forall k: a_k \in D$. Note that $a_3 a_2 a_1 a_0$ represents a sequence of digits, not [multiplication](#).

Notation

When describing base in [mathematical notation](#), the letter b is generally used as a [symbol](#) for this concept, so, for a [binary](#) system, b equals 2. Another common way of expressing the base is writing it as a **decimal** subscript after the number that is being represented (this notation is used in this article). 1111011_2 implies that the number 1111011 is a base-2 number, equal to 123_{10} (a [decimal notation](#) representation), 173_8 ([octal](#)) and 7B_{16} ([hexadecimal](#)). In books and articles, when using initially the written abbreviations of number bases, the base is not subsequently printed: it is assumed that binary 1111011 is the same as 1111011_2 .

The base b may also be indicated by the phrase "base- b ". So binary numbers are "base-2"; octal numbers are "base-8"; decimal numbers are "base-10"; and so on.

To a given radix b the set of digits $\{0, 1, \dots, b-2, b-1\}$ is called the standard set of digits. Thus, binary numbers have digits $\{0, 1\}$; decimal numbers have digits $\{0, 1, 2, \dots, 8, 9\}$; and so on.

Therefore, the following are notational errors: 52_2 , 2_2 , $1A_9$. (In all cases, one or more digits is not in the set of allowed digits for the given base.)

Exponentiation

Positional numeral systems work using **exponentiation** of the base. A digit's value is the digit multiplied by the value of its place. Place values are the number of the base raised to the n th power, where n is the number of other digits between a given digit and the **radix point**. If a given digit is on the left hand side of the radix point (i.e. its value is an **integer**) then n is positive or zero; if the digit is on the right hand side of the radix point (i.e., its value is fractional) then n is negative.

As an example of usage, the number 465 in its respective base b (which must be at least base 7 because the highest digit in it is 6) is equal to:

$$4 \times b^2 + 6 \times b^1 + 5 \times b^0$$

If the number 465 was in base-10, then it would equal:

$$4 \times 10^2 + 6 \times 10^1 + 5 \times 10^0 = 4 \times 100 + 6 \times 10 + 5 \times 1 = 465$$

$$(465_{10} = 465_{10})$$

If however, the number were in base 7, then it would equal:

$$4 \times 7^2 + 6 \times 7^1 + 5 \times 7^0 = 4 \times 49 + 6 \times 7 + 5 \times 1 = 243$$

$$(465_7 = 243_{10})$$

$10_b = b$ for any base b , since $10_b = 1 \times b^1 + 0 \times b^0$. For example, $10_2 = 2$; $10_3 = 3$; $10_{16} = 16_{10}$. Note that the last "16" is indicated to be in base 10. The base makes no difference for one-digit numerals.

This concept can be demonstrated using a diagram. One object represents one unit. When the number of objects is equal to or greater than the base b , then a group of objects is created with b objects. When the number of these groups exceeds b , then a group of these groups of objects is created with b groups of b objects; and so on. Thus the same number in different bases will have different values:

241 in base 5:

2 groups of 5^2 (25)

4 groups of 5

1 group of

1

ooooo ooooo

o o o o o	o o o o o		o o o o o	o o o o o		
o o o o o	o o o o o	+			+	o
o o o o o	o o o o o		o o o o o	o o o o o		
o o o o o	o o o o o					

241 in base 8:

2 groups of 8^2 (64)		4 groups of 8		1 group of 1
o o o o o o o o	o o o o o o o o			
o o o o o o o o	o o o o o o o o			
o o o o o o o o	o o o o o o o o	+	o o o o o o o o	o o o o o o o o
o o o o o o o o	o o o o o o o o			o
o o o o o o o o	o o o o o o o o		o o o o o o o o	o o o o o o o o
o o o o o o o o	o o o o o o o o			
o o o o o o o o	o o o o o o o o			

The notation can be further augmented by allowing a leading minus sign. This allows the representation of negative numbers. For a given base, every representation corresponds to exactly one [real number](#) and every real number has at least one representation. The representations of rational numbers are those representations that are finite, use the bar notation, or end with an infinitely repeating cycle of digits.

Digits and numerals

A *digit* is a symbol that is used for positional notation, and a *numeral* consists of one or more digits used for representing a [number](#) with positional notation. Today's most common digits are the [decimal digits](#) "0", "1", "2", "3", "4", "5", "6", "7", "8", and "9". The distinction between a digit and a numeral is most pronounced in the context of a number base.

A non-zero *numeral* with more than one digit position will mean a different number in a different number base, but in general, the *digits* will mean the same.^[12] For example, the base-8 numeral 23_8 contains two digits, "2" and "3", and with a base number (subscripted) "8". When converted to base-10, the 23_8 is equivalent to 19_{10} , i.e. $23_8 = 19_{10}$. In our notation here, the subscript "8" of the numeral 23_8 is part of the numeral, but this may not always be the case.

Imagine the numeral "23" as having [an ambiguous base](#) number. Then "23" could likely be any base, from base-4 up. In base-4, the "23" means 11_{10} , i.e. $23_4 = 11_{10}$. In base-60, the "23" means the number 123_{10} , i.e. $23_{60} = 123_{10}$. The numeral "23" then, in this case, corresponds to the set of base-10 numbers {11, 13, 15, 17, 19, 21, **23**, ..., 121, 123} while its digits "2" and "3" always retain their original meaning: the "2" means "two of", and the "3" means "three of".

In certain applications when a numeral with a fixed number of positions needs to represent a greater number, a higher number-base with more digits per position can be used. A three-digit, decimal numeral can represent only up to **999**. But if the number-base is increased to 11, say, by adding the digit "A", then the same three positions, maximized to "AAA", can represent a number as great as **1330**. We could increase the number base again and assign "B" to 11, and so on (but there is also a possible encryption between number and digit in the number-digit-numeral hierarchy). A three-digit numeral "ZZZ" in base-60 could mean **215 999**. If we use the entire collection of our [alphanumerics](#) we could ultimately serve a base-62 numeral system, but we remove two digits, uppercase "I" and uppercase "O", to reduce confusion with digits "1" and "0".^[13] We are left with a base-60, or sexagesimal numeral system utilizing 60 of the 62 standard alphanumerics. (But see [Sexagesimal system](#) below.) In general, the number of possible values that can be represented by a *d* digit number in base *r* is r^d .

The common numeral systems in computer science are binary (radix 2), octal (radix 8), and hexadecimal (radix 16). In [binary](#) only digits "0" and "1" are in the numerals. In the [octal](#) numerals, are the eight digits 0–7. [Hex](#) is 0–9 A–F, where the ten numerics retain their usual meaning, and the alphabets correspond to values 10–15, for a total of sixteen digits. The numeral "10" is binary numeral "2", octal numeral "8", or hexadecimal numeral "16".

Radix point

The notation can be extended into the negative exponents of the base *b*. Thereby the so-called radix point, mostly ».«, is used as separator of the positions with non-negative from those with negative exponent.

Numbers that are not [integers](#) use places beyond the [radix point](#). For every position behind this point (and thus after the units digit), the exponent *n* of the power b^n decreases by 1 and the power approaches 0. For example, the number 2.35 is equal to:

$$2 \times 10^0 + 3 \times 10^{-1} + 5 \times 10^{-2}$$

Sign

If the base and all the digits in the set of digits are non-negative, negative numbers cannot be expressed. To overcome this, a **minus sign**, here »-«, is added to the numeral system. In the usual notation it is prepended to the string of digits representing the otherwise non-negative number.

Base conversion

The conversion to a base b_2 of an integer n represented in base b_1 can be done by a succession of **Euclidean divisions** by b_2 : the right-most digit in base b_2 is the remainder of the division of n by b_2 ; the second right-most digit is the remainder of the division of the quotient by b_2 , and so on. The left-most digit is the last quotient. In general, the k th digit from the right is the remainder of the division by b_2 of the $(k-1)$ th quotient.

For example: converting $A10B_{Hex}$ to decimal (41227):

```
0xA10B/10 = 0x101A R: 7 (ones place)
0x101A/10 = 0x19C R: 2 (tens place)
0x19C/10 = 0x29 R: 2 (hundreds place)
0x29/10 = 0x4 R: 1 ...
4
```

When converting to a larger base (such as from binary to decimal), the remainder represents b_2 as a single digit, using digits from b_1 . For example: converting $0b11111001$ (binary) to 249 (decimal):

```
0b11111001/10 = 0b11000 R: 0b1001 (0b1001 = "9" for ones place)
0b11000/10 = 0b10 R: 0b100 (0b100 = "4" for tens)
0b10/10 = 0b0 R: 0b10 (0b10 = "2" for hundreds)
```

For the **fractional** part, conversion can be done by taking digits after the radix point (the numerator), and **dividing** it by the **implied denominator** in the target radix. Approximation may be needed due to a possibility of **non-terminating digits** if the **reduced** fraction's denominator has a prime factor other than any of the base's prime factor(s) to convert to. For example, 0.1 in decimal ($1/10$) is $0b1/0b1010$ in binary, by dividing this in that radix, the result is $0b0.\overline{00011}$

(because one of the prime factors of 10 is 5). For more general fractions and bases see the [algorithm for positive bases](#).

In practice, [Horner's method](#) is more efficient than the repeated division required above^[14]. A number in positional notation can be thought of as a polynomial, where each digit is a coefficient. Coefficients can be larger than one digit, so an efficient way to convert bases is to convert each digit, then evaluate the polynomial via Horner's method within the target base. Converting each digit is a simple [lookup table](#), removing the need for expensive division or modulus operations; and multiplication by x becomes right-shifting. However, other polynomial evaluation algorithms would work as well, like [repeated squaring](#) for single or sparse digits.

Example:

```
Convert 0xA10B to 41227
```

$$A10B = (10*16^3) + (1*16^2) + (0*16^1) + (11*16^0)$$

```
Lookup table:
```

```
0x0 = 0
```

```
0x1 = 1
```

```
...
```

```
0x9 = 9
```

```
0xA = 10
```

```
0xB = 11
```

```
0xC = 12
```

```
0xD = 13
```

```
0xE = 14
```

```
0xF = 15
```

```
Therefore 0xA10B's decimal digits are 10, 1, 0, and 11.
```

```
Lay out the digits out like this. The most significant digit (10)
is "dropped":
```

```
10 1 0 11 <- Digits of 0xA10B
```

```
-----
```

```
10
```

```
Then we multiply the bottom number from the source base (16), the
product is placed under the next digit of the source value, and
```

then add:

```
10 1 0 11
    160
-----
```

```
10 161
```

Repeat until the final addition is performed:

```
10 1 0 11
    160 2576 41216
-----
```

```
10 161 2576 41227
```

and that is 41227 in decimal.

Convert 0b11111001 to 249

Lookup table:

0b0 = 0

0b1 = 1

Result:

```
1 1 1 1 1 0 0 1 <- Digits of 0b11111001
```

```
2 6 14 30 62 124 248
```

```
-----
1 3 7 15 31 62 124 249
```

Terminating fractions

The numbers which have a finite representation form the [semiring](#)

$$\frac{\mathbb{N}_0}{b^{\mathbb{N}_0}} := \{mb^{-\nu} \mid m \in \mathbb{N}_0 \wedge \nu \in \mathbb{N}_0\}.$$

More explicitly, if $p_1^{\nu_1} \cdot \dots \cdot p_n^{\nu_n} := b$ is a [factorization](#) of b into the primes $p_1, \dots, p_n \in \mathbb{P}$ with exponents $\nu_1, \dots, \nu_n \in \mathbb{N}$,^[15] then with the non-empty set of denominators

$S := \{p_1, \dots, p_n\}$ we have

$$\mathbb{Z}_S := \left\{ x \in \mathbb{Q} \mid \exists \mu_i \in \mathbb{Z} : x \prod_{i=1}^n p_i^{\mu_i} \in \mathbb{Z} \right\} = \mathfrak{b}^{\mathbb{Z}} \mathbb{Z} = \langle S \rangle^{-1} \mathbb{Z}$$

where $\langle S \rangle$ is the group generated by the $p \in S$ and $\langle S \rangle^{-1} \mathbb{Z}$ is the so-called **localization** of \mathbb{Z} with respect to S .

The **denominator** of an element of \mathbb{Z}_S contains if reduced to lowest terms only prime factors out of S . This **ring** of all terminating fractions to base \mathfrak{b} is **dense** in the field of **rational numbers** \mathbb{Q} . Its **completion** for the usual (Archimedean) metric is the same as for \mathbb{Q} , namely the real numbers \mathbb{R} . So, if $S = \{p\}$ then $\mathbb{Z}_{\{p\}}$ has not to be confused with $\mathbb{Z}_{(p)}$, the **discrete valuation ring** for the **prime** p , which is equal to \mathbb{Z}_T with $T = \mathbb{P} \setminus \{p\}$.

If \mathfrak{b} divides \mathfrak{c} , we have $\mathfrak{b}^{\mathbb{Z}} \mathbb{Z} \subseteq \mathfrak{c}^{\mathbb{Z}} \mathbb{Z}$.

Infinite representations

Rational numbers

The representation of non-integers can be extended to allow an infinite string of digits beyond the point. For example, 1.12112111211112 ... base-3 represents the sum of the infinite **series**:

$$\begin{aligned} & 1 \times 3^0 + \\ & 1 \times 3^{-1} + 2 \times 3^{-2} + \\ & 1 \times 3^{-3} + 1 \times 3^{-4} + 2 \times 3^{-5} + \\ & 1 \times 3^{-6} + 1 \times 3^{-7} + 1 \times 3^{-8} + 2 \times 3^{-9} + \\ & 1 \times 3^{-10} + 1 \times 3^{-11} + 1 \times 3^{-12} + 1 \times 3^{-13} + 2 \times 3^{-14} + \dots \end{aligned}$$

Since a complete infinite string of digits cannot be explicitly written, the trailing ellipsis (...) designates the omitted digits, which may or may not follow a pattern of some kind. One common pattern is when a finite sequence of digits repeats infinitely. This is designated by drawing a **vinculum** across the repeating block:

$$2.\overline{42314}_5 = 2.42314314314314314 \dots_5$$

This is the **repeating decimal notation** (to which there does not exist a single universally accepted notation or phrasing). For base 10 it is called a repeating decimal or recurring decimal.

An **irrational number** has an infinite non-repeating representation in all integer bases. Whether a **rational number** has a finite representation or requires an infinite repeating representation

depends on the base. For example, one third can be represented by:

$$0.1_3$$

$$0.\overline{3}_{10} = 0.3333333 \dots_{10}$$

or, with the base implied:

$$0.\overline{3} = 0.3333333 \dots \text{ (see also } 0.999\dots \text{)}$$

$$0.\overline{01}_2 = 0.010101 \dots_2$$

$$0.2_6$$

For integers p and q with $\gcd(p, q) = 1$, the fraction p/q has a finite representation in base b if and only if each prime factor of q is also a prime factor of b .

For a given base, any number that can be represented by a finite number of digits (without using the bar notation) will have multiple representations, including one or two infinite representations:

1. A finite or infinite number of zeroes can be appended:

$$3.46_7 = 3.460_7 = 3.460000_7 = 3.46\overline{0}_7$$

2. The last non-zero digit can be reduced by one and an infinite string of digits, each corresponding to one less than the base, are appended (or replace any following zero digits):

$$3.46_7 = 3.45\overline{6}_7$$

$$1_{10} = 0.\overline{9}_{10} \quad \text{(see also } 0.999\dots \text{)}$$

$$220_5 = 214.\overline{4}_5$$

Irrational numbers

A (real) irrational number has an infinite non-repeating representation in all integer bases.

Examples are the non-solvable n th roots

$$y = \sqrt[n]{x}$$

with $y^n = x$ and $y \notin \mathbf{Q}$, numbers which are called algebraic, or numbers like

$$\pi, e$$

which are transcendental. The number of transcendentals is uncountable and the sole way to write them down with a finite number of symbols is to give them a symbol or a finite sequence of symbols.

Applications

Decimal system

In the **decimal** (base-10) **Hindu–Arabic numeral system**, each position starting from the right is a higher power of 10. The first position represents 10^0 (1), the second position 10^1 (10), the third position 10^2 (10×10 or 100), the fourth position 10^3 ($10 \times 10 \times 10$ or 1000), and so on.

Fractional values are indicated by a **separator**, which can vary in different locations. Usually this separator is a period or **full stop**, or a **comma**. Digits to the right of it are multiplied by 10 raised to a negative power or exponent. The first position to the right of the separator indicates 10^{-1} (0.1), the second position 10^{-2} (0.01), and so on for each successive position.

As an example, the number 2674 in a base-10 numeral system is:

$$(2 \times 10^3) + (6 \times 10^2) + (7 \times 10^1) + (4 \times 10^0)$$

or

$$(2 \times 1000) + (6 \times 100) + (7 \times 10) + (4 \times 1).$$

Sexagesimal system

The **sexagesimal** or base-60 system was used for the integral and fractional portions of **Babylonian numerals** and other Mesopotamian systems, by **Hellenistic** astronomers using **Greek numerals** for the fractional portion only, and is still used for modern time and angles, but only for minutes and seconds. However, not all of these uses were positional.

Modern time separates each position by a colon or a **prime symbol**. For example, the time might be 10:25:59 (10 hours 25 minutes 59 seconds). Angles use similar notation. For example, an angle might be $10^\circ 25' 59''$ (10 **degrees** 25 **minutes** 59 **seconds**). In both cases, only minutes and seconds use sexagesimal notation—angular degrees can be larger than 59 (one rotation around a circle is 360° , two rotations are 720° , etc.), and both time and angles use decimal fractions of a second. This contrasts with the numbers used by Hellenistic and **Renaissance** astronomers, who used **thirds**, **fourths**, etc. for finer increments. Where we might write $10^\circ 25' 59.392''$, they would have written $10^\circ 25' 59'' 23''' 31'''' 12'''''$ or $10^\circ 25' 59'' 23''' 31'''' 12'''''$.

Using a digit set of digits with upper and lowercase letters allows short notation for sexagesimal numbers, e.g. 10:25:59 becomes 'ARz' (by omitting I and O, but not i and o), which is useful for use in URLs, etc., but it is not very intelligible to humans.

In the 1930s, [Otto Neugebauer](#) introduced a modern notational system for Babylonian and Hellenistic numbers that substitutes modern decimal notation from 0 to 59 in each position, while using a semicolon (;) to separate the integral and fractional portions of the number and using a comma (,) to separate the positions within each portion.^[16] For example, the mean [synodic month](#) used by both Babylonian and Hellenistic astronomers and still used in the [Hebrew calendar](#) is 29;31,50,8,20 days, and the angle used in the example above would be written 10;25,59,23,31,12 degrees.

Computing

In [computing](#), the [binary](#) (base-2), octal (base-8) and [hexadecimal](#) (base-16) bases are most commonly used. Computers, at the most basic level, deal only with sequences of conventional zeroes and ones, thus it is easier in this sense to deal with powers of two. The hexadecimal system is used as "shorthand" for binary—every 4 binary digits (bits) relate to one and only one hexadecimal digit. In hexadecimal, the six digits after 9 are denoted by A, B, C, D, E, and F (and sometimes a, b, c, d, e, and f).

The [octal](#) numbering system is also used as another way to represent binary numbers. In this case the base is 8 and therefore only digits 0, 1, 2, 3, 4, 5, 6, and 7 are used. When converting from binary to octal every 3 bits relate to one and only one octal digit.

Hexadecimal, decimal, octal, and a wide variety of other bases have been used for [binary-to-text encoding](#), implementations of [arbitrary-precision arithmetic](#), and other applications.

For a list of bases and their applications, see [list of numeral systems](#).

Other bases in human language

Base-12 systems ([duodecimal](#) or dozenal) have been popular because multiplication and division are easier than in base-10, with addition and subtraction being just as easy. Twelve is a useful base because it has many [factors](#). It is the smallest common multiple of one, two, three, four and six. There is still a special word for "dozen" in English, and by analogy with the word for 10^2 , *hundred*, commerce developed a word for 12^2 , *gross*. The standard 12-hour clock and common use of 12 in English units emphasize the utility of the base. In addition, prior to its conversion to decimal, the old British currency [Pound Sterling](#) (GBP) *partially* used base-12; there were 12 pence (d) in a shilling (s), 20 shillings in a pound (£), and therefore 240 pence in a pound. Hence the term LSD or, more properly, [£sd](#).

The [Maya civilization](#) and other civilizations of [pre-Columbian Mesoamerica](#) used base-20 ([vigesimal](#)), as did several North American tribes (two being in southern California). Evidence of base-20 counting systems is also found in the languages of central and western [Africa](#).

Remnants of a [Gaulish](#) base-20 system also exist in French, as seen today in the names of the numbers from 60 through 99. For example, sixty-five is *soixante-cinq* (literally, "sixty [and] five"), while seventy-five is *soixante-quinze* (literally, "sixty [and] fifteen"). Furthermore, for any number between 80 and 99, the "tens-column" number is expressed as a multiple of twenty. For example, eighty-two is *quatre-vingt-deux* (literally, four twenty[s] [and] two), while ninety-two is *quatre-vingt-douze* (literally, four twenty[s] [and] twelve). In Old French, forty was expressed as two twenties and sixty was three twenties, so that fifty-three was expressed as two twenties [and] thirteen, and so on.

In English the same base-20 counting appears in the use of "[scores](#)". Although mostly historical, it is occasionally used colloquially. Verse 10 of Psalm 90 in the King James Version of the Bible starts: "The days of our years are threescore years and ten; and if by reason of strength they be fourscore years, yet is their strength labour and sorrow". The Gettysburg Address starts: "Four score and seven years ago".

The [Irish language](#) also used base-20 in the past, twenty being *fichead*, forty *dhá fhichead*, sixty *trí fhichead* and eighty *ceithre fhichead*. A remnant of this system may be seen in the modern word for 40, *daichead*.

The [Welsh language](#) continues to use a [base-20 counting system](#), particularly for the age of people, dates and in common phrases. 15 is also important, with 16–19 being "one on 15", "two on 15" etc. 18 is normally "two nines". A decimal system is commonly used.

The [Inuit languages](#) use a [base-20](#) counting system. Students from [Kaktovik, Alaska](#) invented a [base-20 numeral system](#) in 1994^[17]

[Danish numerals](#) display a similar [base-20](#) structure.

The [Māori language](#) of New Zealand also has evidence of an underlying base-20 system as seen in the terms *Te Hokowhitu a Tu* referring to a war party (literally "the seven 20s of Tu") and *Tama-hokotahi*, referring to a great warrior ("the one man equal to 20").

The [binary system](#) was used in the Egyptian Old Kingdom, 3000 BC to 2050 BC. It was cursive by rounding off rational numbers smaller than 1 to $1/2 + 1/4 + 1/8 + 1/16 + 1/32 + 1/64$, with a $1/64$ term thrown away (the system was called the [Eye of Horus](#)).

A number of [Australian Aboriginal languages](#) employ binary or binary-like counting systems. For example, in [Kala Lagaw Ya](#), the numbers one through six are *urapon*, *ukasar*, *ukasar-urapon*, *ukasar-ukasar*, *ukasar-ukasar-urapon*, *ukasar-ukasar-ukasar*.

North and Central American natives used base-4 ([quaternary](#)) to represent the four cardinal directions. Mesoamericans tended to add a second base-5 system to create a modified base-20 system.

A base-5 system ([quinary](#)) has been used in many cultures for counting. Plainly it is based on the number of digits on a human hand. It may also be regarded as a sub-base of other bases, such as base-10, base-20, and base-60.

A base-8 system ([octal](#)) was devised by the [Yuki tribe](#) of Northern California, who used the spaces between the fingers to count, corresponding to the digits one through eight.^[18] There is also linguistic evidence which suggests that the Bronze Age [Proto-Indo Europeans](#) (from whom most European and Indic languages descend) might have replaced a base-8 system (or a system which could only count up to 8) with a base-10 system. The evidence is that the word for 9, *newm*, is suggested by some to derive from the word for "new", *newo-*, suggesting that the number 9 had been recently invented and called the "new number".^[19]

Many ancient counting systems use five as a primary base, almost surely coming from the number of fingers on a person's hand. Often these systems are supplemented with a secondary base, sometimes ten, sometimes twenty. In some [African languages](#) the word for five is the same as "hand" or "fist" ([Dyola language](#) of [Guinea-Bissau](#), [Banda language](#) of [Central Africa](#)). Counting continues by adding 1, 2, 3, or 4 to combinations of 5, until the secondary base is reached. In the case of twenty, this word often means "man complete". This system is referred to as *quinquagesimal*. It is found in many languages of the [Sudan](#) region.

The [Telefol language](#), spoken in [Papua New Guinea](#), is notable for possessing a base-27 numeral system.

Non-standard positional numeral systems

Interesting properties exist when the base is not fixed or positive and when the digit symbol sets denote negative values. There are many more variations. These systems are of practical and theoretic value to computer scientists.

Balanced ternary^[20] uses a base of 3 but the digit set is $\{\bar{1}, 0, 1\}$ instead of $\{0, 1, 2\}$. The " $\bar{1}$ " has an equivalent value of -1 . The negation of a number is easily formed by switching the $\bar{1}$ on the 1s. This system can be used to solve the **balance problem**, which requires finding a minimal set of known counter-weights to determine an unknown weight. Weights of 1, 3, 9, ... 3^n known units can be used to determine any unknown weight up to $1 + 3 + \dots + 3^n$ units. A weight can be used on either side of the balance or not at all. Weights used on the balance pan with the unknown weight are designated with $\bar{1}$, with 1 if used on the empty pan, and with 0 if not used. If an unknown weight W is balanced with 3 (3^1) on its pan and 1 and 27 (3^0 and 3^3) on the other, then its weight in decimal is 25 or $10\bar{1}1$ in balanced base-3.

$$10\bar{1}1_3 = 1 \times 3^3 + 0 \times 3^2 - 1 \times 3^1 + 1 \times 3^0 = 25.$$

The **factorial number system** uses a varying radix, giving **factorials** as place values; they are related to **Chinese remainder theorem** and **residue number system** enumerations. This system effectively enumerates permutations. A derivative of this uses the **Towers of Hanoi** puzzle configuration as a counting system. The configuration of the towers can be put into 1-to-1 correspondence with the decimal count of the step at which the configuration occurs and vice versa.

Decimal equivalents	-3	-2	-1	0	1	2	3	4	5	6	7	8
Balanced base 3	$\bar{1}0$	$\bar{1}1$	$\bar{1}$	0	1	$1\bar{1}$	10	11	$1\bar{1}\bar{1}$	$1\bar{1}0$	$1\bar{1}1$	$10\bar{1}$
Base -2	1101	10	11	0	1	110	111	100	101	11010	11011	11000
Factoroid				0	10	100	110	200	210	1000	1010	1100

Non-positional positions

Each position does not need to be positional itself. Babylonian sexagesimal numerals were positional, but in each position were groups of two kinds of wedges representing ones and tens (a narrow vertical wedge (|) and an open left pointing wedge (<))—up to 14 symbols per position (5 tens (<<<<<) and 9 ones (|||||)) grouped into one or two near squares containing up to three tiers of symbols, or a place holder (\\) for the lack of a position).^[21] Hellenistic astronomers used one or two alphabetic Greek numerals for each position (one chosen from 5 letters representing 10–50 and/or one chosen from 9 letters representing 1–9, or a **zero symbol**).^[22]

See also

Examples:

- [List of numeral systems](#)
- [Category: Positional numeral systems](#)

Related topics:

- [Algorism](#)
- [Hindu–Arabic numeral system](#)
- [Mixed radix](#)
- [Non-standard positional numeral systems](#)
- [Numeral system](#)
- [Scientific notation](#)

Other:

- [Significant figures](#)

Notes

1. *Kaplan, Robert (2000). [The Nothing That Is: A Natural History of Zero](https://archive.org/details/nothing_thatisnat00kapl) (https://archive.org/details/nothing_thatisnat00kapl) . Oxford: Oxford University Press. pp. 11–12 – via archive.org.*
2. *"Greek numerals" (https://web.archive.org/web/20161126013536/http://www-gap.dcs.st-and.ac.uk/~history/HistTopics/Greek_numbers.html) . Archived from the original (http://www-gap.dcs.st-and.ac.uk/~history/HistTopics/Greek_numbers.html) on 26 November 2016. Retrieved 31 May 2016.*
3. *Menninger, Karl: Zahlwort und Ziffer. Eine Kulturgeschichte der Zahl, Vandenhoeck und Ruprecht, 3rd. ed., 1979, ISBN 3-525-40725-4, pp. 150–153*
4. *Ibrah, page 187*
5. *L. F. Menabrea. Translated by Ada Augusta, Countess of Lovelace. "Sketch of The Analytical Engine Invented by Charles Babbage" (http://www.fourmilab.ch/babbage/sketch.html) Archived (https://web.archive.org/web/20080915134651/http://www.fourmilab.ch/babbage/sketch.html) 15 September 2008 at the Wayback Machine. 1842.*
6. *Berggren, J. Lennart (2007). "Mathematics in Medieval Islam". The Mathematics of Egypt, Mesopotamia, China, India, and Islam: A Sourcebook. Princeton University Press. p. 518. ISBN 978-0-691-11485-9.*
7. *Gandz, S.: The invention of the decimal fractions and the application of the exponential calculus by Immanuel Bonfils of Tarascon (c. 1350), Isis 25 (1936), 16–45.*

8. [Lam Lay Yong](#), "The Development of Hindu-Arabic and Traditional Chinese Arithmetic", *Chinese Science*, 1996 p38, Kurt Vogel notation
9. [Lay Yong, Lam](#). "A Chinese Genesis, Rewriting the history of our numeral system". *Archive for History of Exact Sciences*. **38**: 101–108.
10. [B. L. van der Waerden](#) (1985). *A History of Algebra. From Khwarizmi to Emmy Noether* (<https://archive.org/details/historyofalgebra0000waer>) . Berlin: Springer-Verlag.
11. [E. J. Dijksterhuis](#) (1970) *Simon Stevin: Science in the Netherlands around 1600*, [Martinus Nijhoff Publishers](#), Dutch original 1943
12. The digit will retain its meaning in other number bases, in general, because a higher number base would normally be a notational extension of the lower number base in any systematic organization. In the [mathematical sciences](#) there is virtually only one positional-notation numeral system for each base below 10, and this extends with few, if insignificant, variations on the choice of alphabetic digits for those bases above 10.
13. We do not usually remove the lowercase digits "l" and lowercase "o", for in most fonts they are discernible from the digits "1" and "0".
14. User 'Gone'. "number systems - How to change from base n to m " (<https://math.stackexchange.com/a/50122>) . Mathematics Stack Exchange. Retrieved 6 August 2020. {{cite web}}: |last1= has generic name (help)
15. The exact size of the ν_1, \dots, ν_n does not matter. They only have to be ≥ 1 .
16. [Neugebauer, Otto](#); [Sachs, Abraham Joseph](#); [Götze, Albrecht](#) (1945), *Mathematical Cuneiform Texts* (<http://books.google.com/books?id=i-juAAAAMAAJ&q=%22sexagesimal+notation%22&pg=PA1>) , American Oriental Series, vol. 29, New Haven: American Oriental Society and the American Schools of Oriental Research, p. 2, [archived](#) (https://web.archive.org/web/20161001222131/https://books.google.com/books?hl=en&lr=&id=i-juAAAAMAAJ&oi=fnd&pg=PA1&dq=%22sexagesimal+notation%22&ots=nkmW8KtNq_&sig=U_K4iOoKy5Xf70UrbjoTyS3hN2A#v=onepage&q=%22sexagesimal%20notation%22&f=false) from the original on 1 October 2016, retrieved 18 September 2019
17. [Bartley, Wm. Clark](#) (January–February 1997). "Making the Old Way Count" (<http://www.ankn.uaf.edu/sop/SOPv2i1.pdf>) (PDF). *Sharing Our Pathways*. **2** (1): 12–13. [Archived](#) (<https://web.archive.org/web/20130625225547/http://ankn.uaf.edu/SOP/SOPv2i1.pdf>) (PDF) from the original on 25 June 2013. Retrieved 27 February 2017.
18. [Barrow, John D.](#) (1992), *Pi in the sky: counting, thinking, and being* (<https://archive.org/details/piinskycouning00barr>) , Clarendon Press, p. 38, ISBN 9780198539568.
19. ([Mallory & Adams](#) 1997) *Encyclopedia of Indo-European Culture*
20. [Knuth](#), pages 195–213

21. Ifrah, pages 326, 379

22. Ifrah, pages 261–264

References

- O'Connor, John; Robertson, Edmund (December 2000). "Babylonian Numerals" (https://web.archive.org/web/20140911192557/http://www-groups.dcs.st-and.ac.uk/~history/HistTopics/Babylonian_numerals.html) . Archived from the original (http://www-groups.dcs.st-and.ac.uk/~history/HistTopics/Babylonian_numerals.html) on 11 September 2014. Retrieved 21 August 2010.
- Kadvany, John (December 2007). "Positional Value and Linguistic Recursion". *Journal of Indian Philosophy*. **35** (5–6): 487–520. doi:10.1007/s10781-007-9025-5 (<https://doi.org/10.1007%2Fs10781-007-9025-5>) . S2CID 52885600 (<https://api.semanticscholar.org/CorpusID:52885600>) .
- Knuth, Donald (1997). *The art of Computer Programming*. Vol. 2. Addison-Wesley. pp. 195–213. ISBN 0-201-89684-2.
- Ifrah, George (2000). *The Universal History of Numbers: From Prehistory to the Invention of the Computer* (https://archive.org/details/unset0000unse_w3q2) . Wiley. ISBN 0-471-37568-3.
- Kroeber, Alfred (1976) [1925]. *Handbook of the Indians of California* (https://books.google.com/books?id=Plqf_OTz4ukC) . Courier Dover Publications. p. 176. ISBN 9780486233680.

External links

Wikimedia Commons has media related to ***Positional numeral systems***.

- [Accurate Base Conversion](https://web.archive.org/web/20170204004954/http://ultrastudio.org/en/MechengburakalkanApplet-1.7.zip) (<https://web.archive.org/web/20170204004954/http://ultrastudio.org/en/MechengburakalkanApplet-1.7.zip>)
- [The Development of Hindu Arabic and Traditional Chinese Arithmetics](https://web.archive.org/web/20160310032143/http://ibrian.net/navon/paper/the_development_of_hindu_arabic_and_traditional_c.pdf?paperid=1247217) (https://web.archive.org/web/20160310032143/http://ibrian.net/navon/paper/the_development_of_hindu_arabic_and_traditional_c.pdf?paperid=1247217)
- [Implementation of Base Conversion](http://www.cut-the-knot.org/recurrence/conversion.shtml) (<http://www.cut-the-knot.org/recurrence/conversion.shtml>) at cut-the-knot
- [Learn to count other bases on your fingers](http://www.intuitor.com/counting/) (<http://www.intuitor.com/counting/>)

- [Online Arbitrary Precision Base Converter \(https://web.archive.org/web/20161109022004/http://thede toolkit.com/tools/base_conversion\)](https://web.archive.org/web/20161109022004/http://thede toolkit.com/tools/base_conversion)

Retrieved from

["https://en.wikipedia.org/w/index.php?title=Positional_notation&oldid=1094236205"](https://en.wikipedia.org/w/index.php?title=Positional_notation&oldid=1094236205)

Last edited 10 days ago by **Writegood**

WIKIPEDIA
